

RNA splicing factors as oncoproteins and tumour suppressors

Heidi Dvinge^{1,2*}, Eunhee Kim^{3*}, Omar Abdel-Wahab^{3,4} and Robert K. Bradley^{1,2}

Abstract | The recent genomic characterization of cancers has revealed recurrent somatic point mutations and copy number changes affecting genes encoding RNA splicing factors. Initial studies of these 'spliceosomal mutations' suggest that the proteins bearing these mutations exhibit altered splice site and/or exon recognition preferences relative to their wild-type counterparts, resulting in cancer-specific mis-splicing. Such changes in the splicing machinery may create novel vulnerabilities in cancer cells that can be therapeutically exploited using compounds that can influence the splicing process. Further studies to dissect the biochemical, genomic and biological effects of spliceosomal mutations are crucial for the development of cancer therapies targeted at these mutations.

Major spliceosome

A ribonucleoprotein complex consisting of five small nuclear RNAs (termed U1, U2, U4, U5 and U6), each in complex with a set of proteins to form small nuclear ribonucleoprotein complexes (snRNPs), that together are responsible for excision of most introns.

¹Computational Biology Program, Public Health Sciences Division, Fred Hutchinson Cancer Research Center.

²Basic Sciences Division, Fred Hutchinson Cancer Research Center, Seattle, Washington 98109, USA.

³Human Oncology and Pathogenesis Program, Memorial Sloan Kettering Cancer Center.

⁴Leukemia Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, New York 10065, USA.

*These authors contributed equally to this work.

Correspondence to R.K.B. and O.A.-W. rbradley@fredhutch.org; abdelwao@mskcc.org

doi:10.1038/nrc.2016.51
Published online 10 Jun 2016

Splicing of mRNA precursors is required for the maturation of almost all human mRNAs, and furthermore is a key step in the regulation of expression of many genes. Alternative splicing, whereby a pre-mRNA can be processed into different mature mRNAs via splice site selection, enables multiple potential protein products to be generated from a single gene and thereby expands the cellular proteome. Although the functional roles of most isoforms generated by alternative splicing are unknown, specific isoforms have been identified that are selected in cancer owing to their ability to promote neoplastic transformation, cancer progression and/or therapeutic resistance (reviewed in REFS 1,2). In some cases, specific splicing changes are promoted or repressed by recurrent somatic point mutations near splice sites, which may promote cancer by inducing mis-splicing of genes that encode tumour suppressors^{3,4}.

In addition to differential splicing of specific genes, cancers may also exhibit global splicing abnormalities. Genome-wide analyses of cancer transcriptomes^{5–7} have revealed widespread splicing alterations such as inefficient intron removal in neoplastic tissues relative to their normal counterparts, but the functional consequences of these differences are unknown.

Finally, recurrent somatic mutations in genes encoding core spliceosomal proteins and associated RNA splicing factors are present at high frequencies in several cancers^{8–12}. These mutations provide a direct genetic link between dysfunction of the splicing machinery and cancer. Both the genetic spectrum of mutations and functional studies of their consequences indicate that RNA splicing factors can act as proto-oncoproteins and tumour suppressors.

In this Review, we outline the current genetic and functional links between dysregulated and/or mutated RNA splicing factors and cancer. We discuss how recurrent mutations affecting splicing factors might promote the development and/or maintenance of cancer. We describe the challenges inherent in connecting mutations in spliceosomal proteins to specific downstream splicing changes, as well as the importance of testing whether mutated splicing factors dysregulate biological processes other than splicing itself. Finally, we discuss how determining the mechanistic consequences of mutated splicing factors may facilitate the identification of novel therapeutic opportunities to selectively target cancers with spliceosomal mutations.

RNA splicing catalysis and regulation

The following is a simplified summary of the catalysis and regulation of RNA splicing. RNA splicing is essential for processing pre-mRNA transcribed from the more than 90% of human protein-coding genes that contain more than one exon into mature mRNAs before translation into proteins^{13,14}. The primary function of splicing is the removal of non-coding introns, a process carried out by the large macromolecular machineries known as the major spliceosome and the minor spliceosome (reviewed in REF. 15). The major spliceosome consists of five small nuclear ribonucleoprotein complexes (snRNPs, pronounced 'snurps'), U1, U2, U4, U5 and U6, and it is responsible for the excision of more than 99% of human introns. The minor spliceosome contains the U5 snRNP, along with the U11, U12, U4atac and U6atac snRNPs, which are the functional analogues of the corresponding snRNPs in the major spliceosome (reviewed in REF. 16).




Figure 1 | Simplified model of constitutive and alternative splicing. **a** | The key sequence features that govern splicing are shown, including consensus sequences of the 5' and 3' splice sites and sequence motifs bound by *trans*-acting splicing factors (serine/arginine-rich (SR) proteins, heterogeneous nuclear ribonucleoproteins (hnRNPs) and others). Sequence elements required for assembly of the spliceosome onto the pre-mRNA, including the splice sites themselves, polypyrimidine (poly(Y)) tract and branch point, frequently follow the illustrated consensus motifs, whereas the sequences of enhancer or silencer elements depend upon the specific RNA-binding protein that recognizes them. Consensus motifs are illustrated as sequence logos, in which the height of each nucleotide corresponds to its approximate genome-wide frequency in bits. Sequence motifs are illustrated as genomic DNA sequence rather than pre-mRNA sequence (T instead of U). Note that splicing factors such as SR proteins and hnRNPs frequently have context-dependent regulatory roles. **b** | Simplified schematic of intron excision and ligation of two adjacent exons. The steps shown are: recognition of the 5' and 3' splice sites by the U1 and U2 small nuclear ribonucleoprotein complexes (snRNPs), assembly of the snRNPs into the active spliceosome, excision of the intron lariat and ligation of the two exons. **c** | Schematic of constitutive and alternative splicing events. Light blue: constitutive sequence that always forms part of the mature mRNA; mid-blue or dark blue: alternative sequence that can be either included or excluded in the mature mRNA.

Minor spliceosome

A ribonucleoprotein complex that catalyses splicing of a small subset of U12-type introns. The introns recognized by the minor spliceosome are typically defined by sequence elements different from those that define U2-type introns, which are recognized by the major spliceosome.

Small nuclear ribonucleoprotein complexes (snRNPs)

These complexes assemble on pre-mRNA to catalyse splicing.

U2AF complex

A heterodimeric protein complex consisting of U2 small nuclear RNA auxiliary factor 1 (U2AF1) and U2AF2. U2AF2 and U2AF1 bind to the polypyrimidine tract and AG dinucleotide of the 3' splice site to facilitate splice site recognition. Only a subset of AG-dependent 3' splice sites require U2AF1 binding for efficient splice site recognition.

Constitutive splice sites

Splice sites that are always recognized and used by the spliceosome. Similarly, constitutive exons are always included in the mature mRNA.


Each constituent snRNP contains a different short non-coding RNA, an Sm or Sm-like protein complex that is required for the formation of the mature snRNP complex and proteins specific to each snRNP (reviewed in REFS 15, 17).

Intron excision is facilitated by short sequence motifs in the pre-mRNA, in particular at boundaries between the upstream exon and intron (the 5' splice site) and the intron and downstream exon (the 3' splice site) (FIG. 1a). Although splicing itself is catalysed by RNA¹⁸, the proper recognition of splice sites relies on RNA–RNA, RNA–protein and protein–protein interactions. U1 snRNP recognizes and binds to the 5' splice site, whereas U2 snRNP interacts with the branch site region adjacent to the 3' splice site, facilitated by interactions with U2 auxiliary factors (U2AFs, which form the U2AF complex) that bind to the 3' splice site. Following recruitment of the U4/U6.U5 tri-snRNP, the assembled spliceosome enters its active conformation and splicing proceeds via two sequential transesterification reactions (reviewed in REFS 15, 17) (FIG. 1b).

Splice sites are typically categorized as *constitutive* splice sites or *alternative* splice sites, depending on whether they are always (constitutive) or only sometimes (alternative) recognized as splice sites and spliced into the mature mRNA. Splicing of both categories of splice site is catalysed by the same molecular machinery, although the efficient recruitment of spliceosomal proteins to alternative splice sites frequently depends on the binding of additional *trans*-acting factors. Although most splicing reactions in some eukaryotes such as *Saccharomyces cerevisiae* are constitutive — meaning that the same 5' and 3' splice sites are always ligated together — splicing is more complex in mammals.

Transcripts from almost all human multi-exon genes are alternatively spliced, whereby a particular 5' splice site can be joined to different 3' splice sites (or vice versa), frequently in a regulated fashion^{13,14}. Alternative splicing events can be further classified based on how they affect the exonic structure of the mature mRNA (FIG. 1c). The distinction between constitutive and alternative splicing is empirical, and so can vary over

Box 1 | Methods to determine differential RNA splicing using RNA-seq data



Accurate detection and quantification of differential splicing remain bioinformatic challenges^{177,178}. Important aspects of the methods used to measure splicing are described below and include: the quality and origins of the genomic annotation of splicing events, whether to consider full-length isoforms or single splice sites (or single exons) in isolation and the choice of statistical method to identify differences between samples.

Genome-wide splicing annotations are available from both general-purpose genomic databases^{179–181} and specialized software packages¹⁸². These annotations are frequently based on exon–exon junctions inferred from expressed sequence tags (ESTs) and full-length cDNAs, limiting the annotation to the extent that such sequences are available for a given cell type or organism. For example, there are >2.5-fold more alternative splicing events annotated in humans than in mice¹⁸², which is a product of the more complete human genome annotation as well as increased rates of alternative splicing in human compared with mouse tissues¹⁸³. As comparatively few splicing events have been annotated in many model organisms, detecting unannotated splicing (or novel splicing) with RNA-seq is frequently necessary. Novel splicing detection can involve searching for reads overlapping new combinations of known 5' and 3' splice sites of a gene, detecting alternative 5' or 3' ends of exons that constitute novel splice junctions^{178,184} or conducting *de novo* isoform reconstruction^{185,186} (methods ordered by increasing complexity). These approaches are more computationally intensive than using existing annotations, and experimental validation of novel splicing events is crucial.

Differential splicing can be analysed at the level of differentially expressed full-length isoforms or at the level of single splicing events (for example, the inclusion or exclusion of a particular cassette exon; see panel a of the figure). Full-length isoforms are the biologically relevant entities that encode the final protein products¹⁸⁷, but the exon composition of each isoform must be statistically inferred owing to the short read lengths of most high-throughput sequencing platforms. Conversely, focusing on single splicing events may be beneficial when analysing splicing mechanisms and regulation — for example, the altered motif preferences induced by U2 small nuclear RNA auxiliary factor 1 (U2AF1) and serine/arginine-rich splicing factor 2 (SRSF2) mutations^{71,75,87,97} — or when a particular alternatively spliced region has an important impact on gene function. Many such studies focus on single events and then infer full-length isoforms for downstream functional

studies. In either case, splicing is quantified using a 'percent spliced in' value (PSI, or ψ value) ranging from 0 to 100%^{14,182}, defined as the percentage of all mRNAs that correspond to the isoform or alternatively spliced sequence of interest relative to all transcripts of the parent gene (see panel b of the figure), independent of gene expression. (For simplicity, only junction-spanning reads are illustrated in the computation of ψ .)

Quantifying differences in splicing, whether of full-length isoforms or of single events, also involves statistical choices with biological implications. Fold-change in ψ values can be used to quantify differential splicing, but may not correlate well with biological importance. For example, consider an isoform with ψ values of 0.1% and 1% in two samples. This corresponds to a fold-change of 10, yet the isoform is low abundance relative to other isoforms of that gene in both samples. Therefore, many studies instead measure absolute changes in splicing as $\Delta\psi$, the difference in ψ value between two samples, and apply thresholds to $\Delta\psi$ to identify potentially important changes in splicing^{14,182} (for example, using U2AF1-mutant (U2AF1^{S34F}, shown on the y axis) versus wild-type (U2AF1^{WT}, shown on the x axis) acute myeloid leukaemia (AML) samples, with exons satisfying $\Delta\psi \geq 10\%$ (red) or $\Delta\psi \leq -10\%$ (blue) highlighted; see panel c of the figure). This measure is statistically more robust, but may ignore differential splicing of low-abundance yet important isoforms, such as a novel isoform that confers gain-of-function or dominant-negative activity.

Ultimately, the optimal splicing analysis strategy depends upon the specific aims of a given study and the downstream data interpretation plan. Three common goals of splicing studies are described below. First, many studies seek to identify mechanistic changes in the splicing process itself, such as alterations caused by spliceosomal mutations or global differences between tumour and normal samples^{5,7}. In such cases, meta-analyses created by averaging many individual splicing events typically provide the highest statistical power. Second, a study may focus on specific isoforms of biological importance, such as ligand-independent isoforms of the androgen receptor^{188,189}, in which case quantifying entire isoforms is essential. Third, a study may seek to identify differentially spliced isoforms that contribute to a known biological phenotype, such as impaired haematopoiesis or enhanced cell migration. Although the biologically relevant entities are whole isoforms, combining single-event and whole-isoform analysis is frequently useful given the exploratory nature of such studies.

Alternative splice sites

Splice sites that are variably recognized and used by the spliceosome. Similarly, alternative exons (also known as cassette or skipped exons) are sometimes, but not always, included in the mature mRNA. Recognition of alternative splice sites is frequently cell type specific and may rely upon the binding of additional *trans*-acting factors.

Expressed sequence tag (EST). Portions of cDNA sequences.

Unannotated splicing

Splicing events that have not been previously reported by published studies or genomic databases such as Ensembl, UCSC, Vega and RefSeq.

ψ value

The percentage of all mRNAs transcribed from a gene that correspond to a particular isoform or contain a particular alternatively spliced sequence relative to all transcripts of the parent gene. For example, the ψ value for a cassette exon is the fraction of all mRNAs that contain the cassette exon. The ψ value is independent of gene expression and falls within the range 0–100%.

Serine/arginine-rich proteins (SR proteins). A family of splicing factors that frequently promote splicing, although their action is context dependent. Many of these proteins bind to pre-mRNA in a sequence-specific manner to activate splicing. Some members of the family are implicated in other cellular processes, including mRNA export and translation.

Heterogeneous nuclear ribonucleoproteins (hnRNPs). Many members of this protein family are splicing factors, although they also participate in other diverse RNA metabolic processes. These proteins frequently repress splicing, although their actions are context dependent.

time as additional transcript sequencing data become available. For example, high-throughput sequencing has transformed the study of RNA splicing by enabling the rapid and accurate quantification of genome-wide splicing patterns (BOX 1). This has led to the realization that many splice sites and/or exons that were previously classified as constitutive are actually alternatively spliced (for example, in distinct cell types or in response to particular stimuli).

Alternative splicing is frequently regulated by *trans*-acting splicing factors, which bind to sequence motifs that are associated with the promotion (enhancers) or repression (silencers) of splicing. These motifs lie in both exons and introns, and frequently have the strongest effect when in close proximity to a splice site¹⁹ (FIG. 1a; reviewed in REFS 20,21). Splicing enhancers and silencers are bound by diverse RNA-binding proteins, exemplified by the serine/arginine-rich proteins (SR proteins) and heterogeneous nuclear ribonucleoproteins (hnRNPs; reviewed in REFS 22–25). SR proteins contain one or two copies of an RNA recognition motif (RRM) domain at the amino terminus that provides RNA-binding specificity and a carboxy-terminal arginine/serine-rich (RS) domain that contributes to protein–protein interactions^{26,27}. hnRNPs similarly contain both RNA-binding domains and relatively unstructured domains that probably contribute to protein–protein interactions²⁵.

SR proteins and hnRNPs canonically promote and repress splicing, respectively, in a sequence-specific manner²². However, recent studies have revealed further nuances to these roles. The regulatory consequences of SR protein or hnRNP binding within pre-mRNA is frequently context specific, giving rise to complex regulatory relationships (reviewed in REF. 21). For example, SR proteins can promote or repress splicing, depending on where they bind within a pre-mRNA^{28,29}. Similarly, canonical splicing repressors such as the hnRNP polypyrimidine tract binding protein 1 (PTB; also known as PTBP1), can activate splicing in a context-dependent manner^{30,31}. Some hnRNPs act primarily as splicing repressors, whereas others activate exon inclusion³². SR proteins and hnRNPs regulate splicing in diverse ways, including facilitating recruitment of the U1 or U2 snRNP, occluding a splice site, ‘looping out’ an exon and other mechanisms (reviewed in REFS 20,21).

In addition to the SR proteins and hnRNPs, many other RNA-binding proteins regulate splicing. These include the CUG-BP- and ETR-3-like (CELF) proteins (reviewed in REF. 33), Muscblind-like (MBNL) proteins (reviewed in REF. 34), RBFOX proteins, signal transduction and activation of RNA (STAR) proteins, NOVA proteins³⁵, epithelial splicing regulatory proteins (ESRPs), T cell-restricted intracellular antigen 1 (TIA1) and TIA1-like (TIAL1), and many others (reviewed in REF. 21). Furthermore, alterations in the levels of core spliceosomal components such as the snRNP proteins SmB and SmB' (REF. 36) can regulate splicing. Mass spectrometry studies indicate that the spliceosome is associated with more than 170 proteins³⁷, and computational studies of exon recognition suggest that hundreds of

sequence motifs contribute to the regulation of splicing³⁸. Although the total number of splicing factors is unknown, the above studies suggest that hundreds of proteins may have roles in splicing regulation.

Dysregulation of splicing in cancer

Pro-tumorigenic and antitumorigenic splicing factors.

Just as regulation of alternative splicing has essential roles in cellular growth, differentiation and tissue development, dysregulated splicing can give rise to protein isoforms that contribute to tumour establishment, progression and resistance to therapy. Many studies have found links between the altered expression and/or activity of splicing factors, cancer-associated splicing and transformation (reviewed in REFS 1,2) (TABLE 1). SR proteins, hnRNPs and other splicing factors can act as both oncoproteins and tumour suppressors.

Some SR proteins can act as oncoproteins when overexpressed in the correct cellular context. For example, SR splicing factor 1 (SRSF1; also known as ASF and SF2) is upregulated in cancers, including lung, colon and breast cancer^{39,40}. Modest overexpression of SRSF1 drove the immortalization of mouse fibroblasts and human and mouse mammary epithelial cells^{39,40}. SRSF1 promotes transformation in part by inducing mis-splicing of *MNK2* and ribosomal protein S6 kinase B1 (*RPS6KB1*; also known as *S6K1*) and activating the mTOR pathway, which is required for SRSF1-mediated transformation^{39,41}, and by promoting the expression of BIM (also known as BCL2L11) and bridging integrator 1 (BIN1) protein isoforms without pro-apoptotic functions⁴⁰. A recent study proposed that SRSF3 is also an oncoprotein when overexpressed⁴². Consistent with this, SRSF3 downregulation promoted p53-mediated cellular senescence in part by promoting the expression of the p53 β isoform⁴³. *SRSF6* has also been characterized as a proto-oncogene that is frequently overexpressed in human skin cancer⁴⁴. Transgenic overexpression of *SRSF6* from the collagen type Ia1 (*Col1a1*) locus in mice induced hyperplasia of skin sensitized by shaving or wounding, partially through aberrant alternative splicing of tenascin C (*Tnc*)⁴⁴. *SRSF6* may also act as an oncoprotein in lung and colon cancer⁴⁵.

hnRNPs have been implicated in cancer in both pro-tumorigenic and antitumorigenic capacities. For instance, two studies reported that MYC-mediated upregulation of specific hnRNPs (hnRNP A1, hnRNP A2 and PTB) resulted in the exclusion of exon 9 of pyruvate kinase muscle (*PKM*), thus promoting expression of the cancer-associated embryonic *PKM2* isoform and aerobic glycolysis in glioma^{46,47}. Expression of the constitutively active variant of epidermal growth factor receptor (EGFR), EGFRvIII, in gliomas was shown to upregulate hnRNP A1, which, in turn, contributed to alternative splicing of MYC associated factor X (*MAX*) to produce delta *MAX* and further promote glycolytic gene expression and proliferation⁴⁸. The splicing factor hnRNP A2/B1 also acts in a pro-tumorigenic capacity. hnRNP A2/B1 is overexpressed in gliomas, where it correlates with poor prognosis, and its overexpression transforms cells *in vitro*⁴⁹.

Table 1 | **Unmutated splicing factors that function as proto-oncogenes or tumour suppressors**

Splicing factor	Downstream dysregulated isoforms that are functionally linked to cancer
ESRP1 and ESRP2	Promote an epithelial splicing programme to regulate EMT ^{61,62}
hnRNP A1	Contributes to aerobic glycolysis in cancer by promoting the expression of specific isoforms of pyruvate kinase (<i>PKM2</i> isoform) ^{46,47} and <i>MAX</i> (the delta <i>MAX</i> isoform) ⁴⁸
hnRNP A2	Contributes to aerobic glycolysis in cancer by promoting the expression of a specific isoform of pyruvate kinase (<i>PKM2</i> isoform) ^{46,47} . Increases breast cancer cell invasion by promoting the expression of a specific isoform of <i>TP53INP2</i> (REF. 195)
hnRNP A2/B1	Acts as an oncogenic driver in glioblastoma by regulating splicing of several tumour suppressors and oncogenes, including <i>RON</i> ⁴⁹
hnRNP H	Contributes to survival of gliomas and invasion by promoting the expression of specific isoforms of <i>IG20</i> and <i>RON</i> , respectively ¹⁹⁶
hnRNP K	Serves as a tumour suppressor in leukaemia. Deletion is associated with aberrant p21 and C/EBP expression (although mechanistic links to splicing and gene expression are unclear) ⁵¹
hnRNP M	Contributes to EMT in breast cancer and increases metastasis in mice by promoting the expression of a specific isoform of <i>CD44</i> (<i>CD44s</i>) ¹⁹⁷
PRPF6	Promotes cell proliferation in colon cancer by altering splicing of genes associated with growth regulation, including the kinase gene <i>ZAK</i> ¹⁹⁸
PTB (PTBP1)	Contributes to aerobic glycolysis in cancer by promoting the expression of a specific isoform of pyruvate kinase (<i>PKM2</i> isoform) ^{46,47} . Also promotes the expression of an isoform of the deubiquitylating enzyme-encoding gene <i>USP5</i> (REF. 199) that has been shown to promote glioma cell growth and mobility (although the mechanism underlying this phenotypic association is not resolved)
QKI	Acts as a tumour suppressor by regulating alternative splicing of <i>NUMB</i> in lung cancer cells ⁵³
RBFOX2	Promotes a mesenchymal splicing programme to regulate EMT ⁶²
RBM4	Acts as a tumour suppressor by promoting the pro-apoptotic isoform BCL-X _s of <i>BCL2L1</i> and opposing the pro-tumorigenic effects of SRSF1 on mTOR activation ⁶⁰
RBM5, RBM6 and RBM10	RBM5 modulates apoptosis by regulating alternative splicing of <i>CASP2</i> (REF. 200) and <i>FAS</i> ²⁰¹ . RBM5 or RBM6 depletion has an opposite effect to RBM10 depletion, as these factors antagonistically regulate the alternative splicing of <i>NUMB</i> ^{58,59}
SRSF1	Promotes an isoform of the kinase <i>MNK2</i> that promotes eIF4E phosphorylation independently of MAPK signalling ³⁹ . In the context of breast cancer, SRSF1 overexpression promotes alternative splicing of <i>BIM</i> and <i>BIN1</i> (REF. 40) to promote the expression of isoforms that lack pro-apoptotic functions
SRSF3	Regulates alternative splicing of <i>TP53</i> (REF. 43) such that SRSF3 loss promotes expression of p53 β , an isoform of p53 that promotes p53-mediated senescence
SRSF6	Promotes expression of isoforms of the extracellular matrix protein tenascin C that are characteristic of invasive and metastatic skin cancer ⁴⁴ , contributing to epithelial cell hyperplasia
SRSF10	Promotes cell proliferation and colony formation <i>in vitro</i> and increases tumorigenic capacity of colon cancer cells in mice by inducing expression of a specific isoform of <i>BCLAF1</i> (<i>BCLAF1-L</i>) ²⁰²

BCLAF1, BCL-2-associated transcription factor 1; EMT, epithelial-to-mesenchymal transition; ESRP, epithelial splicing regulatory protein; hnRNP, heterogeneous nuclear ribonucleoprotein; NMD, nonsense-mediated decay; PRPF, pre-mRNA processing factor; PTB, polypyrimidine tract-binding protein 1; QKI, quaking; RBM, RNA-binding motif protein; SRSF, serine/arginine-rich splicing factor.

Conversely, other hnRNPs may act as tumour suppressors. Motivated by the observation that *HNRNPK* (among other genes) lies within a chromosomal locus that is recurrently deleted in acute myeloid leukaemia (AML)⁵⁰, one recent mouse study⁵¹ found that deletion of one allele of *Hnrnpk* resulted in myeloid haematological transformation. However, it remains unknown whether the observed myeloid transformation phenotype is due to changes in splicing or other biological pathways. hnRNP K has been implicated in many biological processes in addition to RNA splicing, including cellular proliferation and cellular senescence through its effects on p53 (REF. 52) and p21 (REF. 51), as well as myeloid differentiation through its effects on CCAAT/enhancer-binding protein- α (C/EBP α) and C/EBP β ⁵¹.

Many splicing factors other than SR proteins and hnRNPs have also been implicated in cancer in pro-tumorigenic as well as antitumorigenic capacities. For example, the STAR family protein quaking (QKI) may act as a tumour suppressor in lung cancer, in which it is commonly downregulated⁵³. QKI overexpression inhibits the proliferation of lung cancer cells *in vitro* and *in vivo*, in part by regulating the alternative splicing of *NUMB*⁵³. Other splicing factors are also linked to cancer by their regulation of *NUMB*. RNA binding motif protein 5 (*RBM5*), *RBM6* and *RBM10*, which encode homologous RNA-binding proteins, are commonly deleted, mutated and/or underexpressed or overexpressed in lung and other cancers^{54–58}. *RBM5* or *RBM6* depletion has an opposite effect to *RBM10* depletion

Acute myeloid leukaemia
A type of cancer characterized by the rapid growth of abnormal white blood cells that accumulate in the bone marrow and interfere with the production of normal blood cells.

on *in vitro* colony formation. This is due partly to the antagonistic functions of these factors in regulating the alternative splicing of *NUMB*⁵⁹. RBM4 is also downregulated in multiple cancers and has been characterized as a tumour suppressor that regulates *BCLX* (also known as *BCL2L1*) splicing and antagonizes the pro-tumour

activity of SRSF1 (REF. 60). The epithelial-to-mesenchymal transition (EMT) is regulated by splicing factors including epithelial splicing regulatory protein 1 (ESRP1; also known as RBM35A) and ESRP2 (also known as RBM35B), which promote epithelial splicing programmes in breast cancer, and RBFOX2, which




Figure 2 | Commonly mutated spliceosomal proteins and their associations with specific cancer types. **a** | The incidence and cancer distribution of the most frequently mutated spliceosomal genes are shown. **b** | The spectrum of mutations that have been identified in U2 small nuclear RNA auxiliary factor 1 (U2AF1), serine/arginine-rich splicing factor 2 (SRSF2), zinc finger, RNA-binding motif and serine/arginine-rich 2 (ZRSR2) and splicing factor 3B, subunit 1 (SF3B1). Mutations shown in bold text occur at hot spots; other illustrated mutations are recurrent but rare. As ZRSR2 mutations do not occur at hot spots, very rare or private mutations are shown as examples. **c** | Schematic illustrating the potential reasons that spliceosomal gene

mutations are mutually exclusive of one another: they either converge on a common downstream target or result in synthetic lethality. **d** | The frequency of specific mutations in SF3B1 across various histological subtypes of cancer. AML, acute myeloid leukaemia; CLL, chronic lymphocytic leukaemia; CMMML, chronic myelomonocytic leukaemia; HD, HEAT domain; PPP1R8, binding site for protein phosphatase 1 regulatory subunit 8; RARS, refractory anaemia with ringed sideroblasts; RCMD-RS, refractory cytopenia with multilineage dysplasia and ringed sideroblasts; RRM, RNA recognition motif; RS, arginine/serine-rich domain; sAML, secondary AML; UHM, U2AF homology motif; Zn, zinc finger domain.

Myelodysplastic syndromes (MDS). A heterogeneous group of clonal disorders of haematopoiesis characterized by an impaired ability to generate mature blood cells as well as aberrant cell morphologies (termed dysplasia).

Chronic lymphocytic leukaemia (CLL). A type of cancer characterized by accumulation of aberrant mature-appearing B lymphocytes.

promotes mesenchymal splicing programmes in breast cancer^{61,62}. MBNL and CELF proteins and hnRNPs have also been implicated in EMT⁶², highlighting the complexity of cancer-associated splicing dysregulation.

As described above, both functional and prognostic data indicate that many splicing factors play pro-tumorigenic or antitumorigenic roles. In a few cases, for example, hnRNP K, RBM5, RBM6 and RBM10, genetic changes such as chromosomal deletions may alter the expression of splicing factors. However, in general, the splicing factors described above are not known to be subject to recurrent, high-frequency mutations in cancer. Although these data suggest that dysregulated expression of splicing factors has important roles in tumour development or progression, thus far there is limited functional evidence that altered levels of splicing factors alone can drive cancer initiation or that altered levels of splicing factors are required for cancer maintenance.

Recurrent mutations in spliceosomal genes in cancer.

The discovery of recurrent somatic mutations in genes encoding core spliceosomal proteins throughout diverse cancer types provided the first genetic evidence directly linking RNA splicing regulation to cancer. These spliceosomal mutations were initially discovered in haematological malignancies, including myelodysplastic syndromes (MDS)^{8–10} and other myeloid neoplasms as well as chronic lymphocytic leukaemia (CLL)^{11,12} (FIG. 2a; TABLE 2). More recently, recurrent spliceosomal mutations have also been found in several solid tumour types, including uveal melanoma (15–20%)^{63,64}, pancreatic ductal adenocarcinoma (4%)⁶⁵, lung adenocarcinoma (3%)⁵⁵ and breast cancers (1.8%)^{66–69}. Most reported spliceosomal mutations are concentrated in four genes: splicing factor 3B, subunit 1 (*SF3B1*), serine/arginine-rich splicing factor 2 (*SRSF2*), U2 small nuclear RNA auxiliary factor 1 (*U2AF1*) and zinc finger, RNA-binding motif and serine/arginine-rich 2 (*ZRSR2*)^{8–10}.

Table 2 | Recurrent mutations affecting splicing factors in cancer and their disease associations

Splicing factor	Disease	Mutated residue*	Frequency (%)	Clinical associations
SF3B1	RARS	K700, K666, H662, R625 and E622	48–75.3	Increased event-free or overall survival, lower risk of transformation to AML ¹⁰
	CMML	K700, K666, R625 and E622	4.3–4.7	—
	MDS	K700, K666, H662, R625 and E622	4.4–7	—
	sAML	K700 and K666	3.7–4.8	—
	AML	K700, K666 and G740	2.6–7.3	—
	CLL	K700, K666, H662 and G742	15	Reduced treatment-free or overall survival, faster disease progression ^{11,12,78}
	Pancreatic cancer	K700 and R625	4	—
	Breast cancer	K700, K666, E622 and N626	1.8	—
	Uveal melanoma	R625 and K666	15–20	Relatively favourable prognosis ^{63,64}
	Bladder cancer	R625 and E902	4	—
U2AF1	RARS	S34 and Q157	5	Less favourable overall survival, higher risk of transformation to AML ⁹
	CMML	—	8–11	—
	MDS	—	10.3–11.6	—
	sAML	—	9.7	—
	AML	—	1.3	—
	Lung	S34	3	Reduced progression-free survival ⁵⁵
SRSF2	RARS	P95	5.5	Unfavourable prognostic factor for AML transformation, reduced overall survival ²⁰³
	CMML	—	28–47	—
	MDS	—	11.6	—
	sAML	—	6.5	—
	AML	—	0.7	—
ZRSR2	RARS	No evident mutational hot spot (presumed loss-of-function)	1.4–10	Higher risk of transformation to AML ²⁰⁴
	CMML	—	8	—
	MDS	—	7.7	—
	sAML	—	1.6	—

AML, acute myeloid leukaemia; CMML, chronic myelomonocytic leukaemia; CLL, chronic lymphocytic leukaemia; MDS, myelodysplastic syndromes; RARS, refractory anaemia with ringed sideroblasts; sAML, secondary AML; SF3B1, splicing factor 3B, subunit 1; SRSF2, serine/arginine-rich splicing factor 2; U2AF1, U2 small nuclear RNA auxiliary factor 1; ZRSR2, zinc finger, RNA-binding motif and serine/arginine-rich 2. *Only residues recurrently affected by somatic mutations are listed.

Several features of spliceosomal gene mutations immediately suggested their potential functional consequences. First, with the exception of *ZRSR2* mutations, these mutations affect highly restricted amino acid residues in an exclusively heterozygous state (FIG. 2b). These data suggest that mutations in most spliceosomal genes probably confer gain or alteration of function, with the exception of *ZRSR2* mutations, which follow a pattern consistent with loss of function. Second, splicing factor mutations are mutually exclusive of one another, which may be due to either functional redundancy or synthetic lethality, possibilities that have not yet been explored in published studies (FIG. 2c).

These genetic observations suggest that *SF3B1*, *SRSF2* and *U2AF1* may be proto-oncogenes, as they are subject to highly specific missense mutations suggestive of gain or alteration of function. In contrast, *ZRSR2* may have a tumour suppressor role, as *ZRSR2* mutations frequently introduce in-frame stop codons or disrupt the reading frame, which probably inactivates the gene and/or protein. Functional evidence supporting pro-oncogenic versus tumour suppressor roles for these four proteins is described below.

Although each of these four proteins has some impact on 3' splice site recognition, it is not clear why they are the targets of recurrent mutations in cancer. The more than 30 other proteins that comprise the U2 snRNP, the U2AF complex or the SR protein family are not recurrently mutated despite having similarly important roles in 3' splice site and exon recognition. The highly specific nature of spliceosomal mutations suggests that these mutations, and presumably not others, alter RNA-protein and/or protein-protein interactions in a manner that promotes a specific set of downstream changes that are crucial for oncogenesis.

Specific splicing factors are most frequently mutated in distinct cancer subtypes (FIG. 2a). For example, *SF3B1* is the only splicing factor that has been identified as a common mutational target in breast cancer^{66–69} (based on cohorts sequenced to date), whereas *U2AF1* is the most commonly mutated factor in non-small-cell lung cancer⁵⁵. Moreover, in diseases such as MDS in which multiple splicing factors are commonly mutated, specific splicing factor mutations associate with distinct MDS subtypes. Mutations in *SF3B1* are highly enriched in refractory anaemia with ringed sideroblasts (RARS), a form of MDS characterized by a typically indolent course, anaemia and the accumulation of erythroid precursor cells with abnormally iron-loaded mitochondria^{8,10}. In contrast, *SRSF2* is the most commonly mutated splicing factor in MDS with a more fulminant course⁷⁰ (FIG. 2a). Currently, the mechanistic basis for this association of different splicing factors with distinct histological subtypes of cancer is not known. Moreover, specific mutated residues in *SF3B1* seem to be associated with distinct diseases (FIG. 2d). For instance, *SF3B1* R625 mutations represent the most common *SF3B1* mutation in uveal melanoma, yet are far less frequent in haematological malignancies^{63,64}. Similarly, *U2AF1* mutations affect both the S34 and Q157 residues in haematopoietic malignancies^{8,9}, but only mutations affecting S34 have been identified in lung cancer⁵⁵.

Knock-in mouse models in which mutated spliceosomal genes are expressed from their endogenous loci have been created for *Srsf2* (two distinct models)^{71,72} and *Sf3b1* (REFS 73,74). A transgenic model of inducible *U2AF1*^{S34F} expression has also been described⁷⁵ (BOX 2). Phenotypic analyses of an *Srsf2*^{P95H} conditional knock-in model⁷¹ and the transgenic *U2AF1*^{S34F} model⁷⁵ revealed that the expression of these mutated proteins in postnatal haematopoietic tissue recapitulated key features of MDS, including leukopenia and increased numbers of haematopoietic stem and myeloid progenitor cells. *Srsf2*^{P95H} knock-in mice developed morphological dysplasia whereas the transgenic *U2AF1*^{S34F} mice did not. However, the transcriptomes of haematopoietic stem or progenitor cells (HSPCs) from each model showed that these cells had the same genome-wide alterations in exonic splicing enhancer (*Srsf2*^{P95H}) and 3' splice site (*U2AF1*^{S34F}) preferences as those that were observed in the cells of patients with these mutations. Additionally, in both the *Srsf2*^{P95H} and *U2AF1*^{S34F} mouse models, there were some genes that were differentially spliced in the mouse tumour cells that were also found in patients' cells, suggesting that the models recapitulate many molecular phenotypes of human disease.

The high frequencies with which *SF3B1*, *SRSF2*, *U2AF1* and *ZRSR2* are subject to specific mutations in cancer suggest that spliceosomal mutations drive tumorigenesis, at least in some cellular contexts, and are not merely passenger mutations. Spliceosomal mutations are likely to occur as both initiating and secondary genetic events, a distinction that has been best studied in liquid neoplasms. The clonal architecture of MDS indicates that *SF3B1* mutations are initiating genetic events⁷⁶. Similar studies of secondary AML revealed that *SRSF2*, *U2AF1* and *ZRSR2* mutations also occur early during the leukaemogenic process⁷⁷. In contrast, even though *SF3B1* is the second most commonly mutated gene in CLL, *SF3B1* mutations occur most frequently in advanced as opposed to early disease, suggesting that they are secondary genetic events that facilitate progression^{11,78}.

Mutations affecting 3' splice site recognition via U2. *SF3B1* is mutated at significant rates in both haematological and solid cancers, including MDS, CLL, uveal melanoma and breast cancer, rendering it the most commonly mutated spliceosomal gene (FIG. 2a). It encodes a core spliceosomal protein that binds upstream of the pre-mRNA branch site in a manner that is likely to be largely sequence independent. *SF3B1* is probably required for the recognition of most 3' splice sites¹⁷. *SF3B1* mutations are concentrated in sequence encoding its HEAT repeat domains (FIG. 2b); however, the normal function of these domains is poorly characterized, rendering it difficult to predict the mechanistic consequences of *SF3B1* mutations. DeBoever *et al.*⁷⁹ recently reported that *SF3B1* mutations were associated with enhanced recognition of cryptic 3' splice sites between the branch point and the normal 3' splice site. While U2 snRNP binding to the branch point normally prevents recognition of AG dinucleotides immediately downstream of the branch point by steric occlusion, DeBoever *et al.*⁷⁹ hypothesized that

SF3B1

A gene encoding a key component of the U2 small nuclear ribonucleoprotein complex (snRNP) that binds upstream of the branch point to facilitate 3' splice site recognition. *SF3B1* is probably required for the splicing of most introns and is the most commonly mutated splicing factor in cancer.

SRSF2

A gene encoding a serine/arginine-rich protein (SR protein) that binds to specific exonic splicing enhancer motifs to promote recognition and inclusion of exons containing these motifs.

ZRSR2

A gene encoding a component of the minor spliceosome that contacts the 3' splice site of specific U12-type introns to promote their excision.

Synthetic lethality

The situation in which two cellular perturbations (for example, two distinct mutations, or a mutation and a particular drug) result in cell death when combined whereas each perturbation alone does not.

Stop codons

UAA, UAG or UGA codons, signalling the end of translation. Also known as termination codons.

Exonic splicing enhancer

A typically short sequence motif in pre-mRNA that is bound by a splicing factor to promote exon recognition and subsequent inclusion of the exon in the mature mRNA. Many serine/arginine-rich (SR) proteins bind exonic splicing enhancers to activate splicing.

Secondary AML

(sAML). Acute myeloid leukaemia that develops following a previous chronic myeloid malignancy such as a myelodysplastic syndrome.

Cryptic 3' splice sites

Potential 3' splice sites that are not normally recognized by the spliceosome. By chance, introns and exons contain many AG dinucleotides that are not used as splice sites. Perturbations such as spliceosomal mutations can cause such 'decoy splice sites' to be incorrectly recognized as authentic splice sites.


SF3B1 mutations prevent this normal steric occlusion, thereby enhancing recognition of cryptic splice sites (FIG. 3a). Darman *et al.*⁸⁰ similarly reported that mutant *SF3B1* enhanced recognition of intron-proximal cryptic 3' splice sites, which frequently involved normally unused upstream branch points. However, the exact mechanism or mechanisms by which mutations might alter *SF3B1* interactions with pre-mRNA, components of the U2 snRNP or other proteins remains unknown. A precise understanding of how mutations alter the role of *SF3B1* in RNA splicing will probably require further studies of normal *SF3B1* function.

The consequences of *SF3B1* mutations may be cell type dependent and/or allele specific, as different *SF3B1* mutations may not be phenotypically equivalent.

SF3B1 mutations in MDS as opposed to CLL constitute initial as opposed to secondary genetic insults and associate with favourable as opposed to poor prognosis, respectively^{10–12,78}. Different *SF3B1* mutations are more enriched in MDS compared with CLL and other cancers (FIG. 2d).

Despite the close association between *SF3B1* mutations and the presence of ringed sideroblasts in MDS, no studies have clearly demonstrated that *SF3B1* mutations induce abnormal iron metabolism. However, the recent report of Darman *et al.*⁸⁰ that ATP-binding cassette subfamily B member 7 (*ABCB7*) is mis-spliced in *SF3B1*-mutant cells hints at possible connections between altered splicing and sideroblastic anaemia. *ABCB7* encodes an iron transporter that is essential for

Box 2 | Models of spliceosomal gene mutations in cancer



The genetic heterogeneity of primary patient samples makes genetic models important tools when studying the roles of spliceosomal mutations in cancer pathogenesis. As the wild-type and mutant alleles of spliceosomal genes are consistently co-expressed at similar levels in patients, mimicking this expression pattern is probably important to create biologically realistic genetic models. Analyses of the haematopoietic transcriptomes of serine/arginine-rich splicing factor 2 (*Srsf2*)^{P95H} conditional knock-in⁷¹ (shown in figure panel a; light pink) and U2 small nuclear RNA auxiliary factor 1 (*U2AF1*)^{S34F} transgenic⁷⁵ (shown in figure panel b; light green) mouse models demonstrated that the mechanistic alterations in exon and splice site recognition induced by these mutations are conserved between human and mouse and validate these mouse models for mechanistic studies of the role of splicing alterations in cancer pathogenesis. These observations are consistent with the deep conservation of *SRSF2* and *U2AF1* across eukaryotes. However, even though the functions of these spliceosomal proteins are conserved, the sets of downstream mis-spliced genes may differ between human and mouse models due to species-specific differences in splicing. Figure panel c: tissue-specific splicing events are more frequently conserved between human (pink) and mouse (blue) than their unregulated counterparts, as is the flanking intronic sequence^{14,190}. This even extends to conservation between the human and fungal genomes for a splicing event that enables autoregulation of *SRSF5* (REF. 140).

Figure panel d: nonetheless, most alternative isoforms are species specific^{183,191,192}, such as the illustrated example of *CD55* (pink, human exons; blue, mouse exons). This situation — in which a subset of regulated isoforms is highly conserved, but most alternative splicing events are likely not to be — is analogous to that observed for transcription factor binding, in which detailed studies of transcription factors active in the liver revealed that most binding sites are species specific^{193,194}. It remains to be determined what fraction of species-specific isoforms contributes to phenotypic differences between species compared with those that constitute non-functional products of noisy splicing. However, as only approximately one-quarter of alternative exons are conserved between human and mouse¹⁸³, mis-spliced isoforms that are relevant to human disease pathogenesis yet absent from the mouse genome are likely to exist. Given these differences between species, the parallel use of genome-engineering techniques to generate isogenic human cell lines carrying spliceosomal gene mutations in endogenous loci may prove particularly useful for detailed transcriptional studies. For example, Zhang *et al.*⁹⁷ recently generated *SRSF2*^{P95H} knock-in K562 cells to identify the changes in exon recognition and differential splicing induced by *SRSF2* mutations. Combined studies of mouse models, isogenic human cells and patient cohorts is likely to prove essential to identify the direct targets of mutant spliceosomal proteins with relevance for cancer. HSPCs, haematopoietic stem and progenitor cells.

haematopoiesis and that is mutated in X-linked sideroblastic anaemia with ataxia, a genetic disease that is also characterized by the presence of ringed sideroblasts^{81,82}.

U2AF1 is mutated in both liquid and solid tumours, with the highest reported rates in MDS and lung cancer (FIG. 2a). Unlike SF3B1, which is required for recognition of many or most 3' splice sites, U2AF1 binds to the 3' splice site in a highly sequence-specific manner, and is required for recognition of only a subset of 'AG-dependent' 3' splice sites^{83,84}. *U2AF1* mutations are concentrated in sequence that encodes the S34 and Q157 residues, which lie within the first and second C₃H zinc 'knuckles' of the protein (FIG. 2b). In haematological cancer, such mutations have been shown to induce sequence-specific alterations in the preferred RNA motif bound by U2AF1, which normally recognizes the motif yAG|r (y = pyrimidine; r = purine; lower-case indicates nucleotides that are preferred but not always required, whereas upper-case indicates nucleotides that are usually required; 'I' = intron–exon boundary)^{85–88} (FIG. 3b). Interestingly, the molecular consequences of *U2AF1* mutations are allele specific⁸⁷. S34 and Q157 mutations affect recognition of the –3 (pyrimidine normally preferred) and +1 (purine normally preferred) positions, respectively, where the coordinates are defined with respect to the intron–exon boundary, to induce different changes in 3' splice site recognition. At a global level, S34 mutations promote recognition of 3' splice sites with cAG|r and aAG|r motifs over those with tAG|r, whereas Q157 mutations promote 3' splice sites with yAG|g motifs over those with yAG|a motifs.

Many downstream targets of mutant *U2AF1* have been identified using patient transcriptomes, a transgenic mouse model of *U2AF1*^{S34F} and transgenic human cells bearing each of the common *U2AF1* mutations that affect the S34 and Q157 residues^{75,85–88}. These mis-spliced genes fall into biological pathways that include the DNA damage response (ataxia telangiectasia and Rad3-related (*ATR*) and Fanconi anaemia complementation group A (*FANCA*)), epigenetic regulation (*H2A* histone family member Y (*H2AFY*)) and apoptosis (caspase 8 (*CASP8*))^{75,87}. Intriguingly, several of the identified target genes are recurrently mutated in MDS (for example, *BCL6* co-repressor (*BCOR*)⁸⁹) and other cancers (for example, *CASP8*; REFS 66,90). However, functional studies are needed to determine whether these and other mis-spliced genes contribute to the disease process.

Mutations affecting 3' splice site recognition via U12. *ZRSR2* mutations found in MDS are distributed throughout the gene, which lies on the X chromosome (Xp22.1), and frequently interrupt the coding sequence by directly or indirectly introducing in-frame stop codons (FIG. 2a). Together with the common occurrence of *ZRSR2* mutations in male patients with cancer⁸, this pattern is consistent with loss of function. *ZRSR2* mutations therefore contrast with the mutations observed in the spliceosomal genes *SF3B1*, *SRSF2* and *U2AF1*, which cause missense changes at specific residues and never introduce in-frame stop codons (FIG. 2b).

Whereas biochemical assays suggested that *ZRSR2* promotes recognition of both U2- and U12-type introns, the phylogenetic observation that organisms with U12-type introns have *ZRSR2* and those lacking U12-type introns also lack *ZRSR2* suggested that *ZRSR2* is particularly important for U12-type splicing⁹¹. Madan *et al.*⁹² reported that MDS transcriptomes harbouring mutations that are likely to inactivate *ZRSR2* are characterized by frequent retention of U12-type introns, consistent with a crucial role for *ZRSR2* in the minor spliceosome (FIG. 3c). *ZRSR2* knockdown altered the *in vitro* differentiation potential of cord blood-derived CD34⁺ cells by promoting myeloid differentiation and impairing erythroid differentiation⁹², consistent with features of human MDS. However, *ZRSR2* knockdown also impaired the growth of K562 cells *in vitro* and following subcutaneous xenograft *in vivo*⁹², indicating that *ZRSR2* loss does not convey a proliferative advantage in the K562 genetic background. *ZRSR2* mutations were associated with mis-splicing of genes relevant to the MAPK pathway and E2F transcription factor signalling⁹², but functional experiments are needed to determine whether these or other splicing changes contribute to the haematopoietic phenotypes of *ZRSR2*-deficient cells.

Mutations affecting exon recognition. *SRSF2* mutations occur most commonly (in 40–50% of patients⁸) in chronic myelomonocytic leukaemia (CMML), and are also enriched in subtypes of high-risk MDS, in which they portend an increased risk of transformation to acute leukaemia⁹³ (FIG. 2a). *SRSF2* encodes a member of the SR protein family that contributes to both constitutive and alternative splicing. *SRSF2* canonically promotes exon recognition by binding to exonic splicing enhancer (ESE) motifs within pre-mRNAs through its RRM domain^{94–96}. All recurrent *SRSF2* mutations affect the P95 residue⁸, which is immediately downstream of the RRM domain (FIG. 2b). RNA-seq analyses of HSPCs from *Srsf2*^{P95H} knock-in mice⁷¹ and transgenic⁷¹ and knock-in⁹⁷ K562 cells expressing *SRSF2*^{P95H/L/R}, as well as human AML and CMML patients with or without *SRSF2* mutations⁷¹, revealed that *SRSF2* mutations alter its normal sequence-specific RNA-binding activity. Mutant *SRSF2* preferentially recognizes a C-rich CCNG motif versus a G-rich GGNG motif, whereas wild-type *SRSF2* binds to both motifs with similar affinity^{71,97,98} (FIG. 3d). These alterations in the RNA-binding activity of *SRSF2* promote or repress recognition of exons containing C-rich or G-rich ESEs^{71,97}.

Altered ESE recognition causes widespread splicing changes in hundreds of genes^{71,97}. Several of these downstream mis-spliced genes are themselves recurrent mutational targets in myeloid malignancies, including enhancer of zeste homologue 2 (*EZH2*)^{99,100} and *BCOR*⁸⁹. *SRSF2* mutations promote inclusion of a 'poison exon' of *EZH2* that introduces an in-frame stop codon to induce nonsense-mediated decay (NMD) of the *EZH2* transcript and consequent global downregulation of *EZH2* protein and histone H3 lysine 27 trimethylation (H3K27me3) levels⁷¹. Loss-of-function mutations in

Chronic myelomonocytic leukaemia

(CMML). A clonal disorder with features of both myelodysplastic and myeloproliferative syndromes in which there are too many monocytes in the blood.

Poison exon

A cassette exon containing an in-frame premature stop codon. A premature stop codon lies upstream of the normal stop codon, resulting in premature termination of translation of the mRNA when it is included in a transcript. Poison exons can induce nonsense-mediated decay of the mRNA or production of a truncated protein.

Nonsense-mediated decay

(NMD). An RNA surveillance process that recognizes and degrades mRNAs containing premature stop codons, as well as other abnormal RNAs and a subset of normal coding transcripts. Splicing is closely linked to NMD, as exon–exon junctions are important components of NMD activation in human cells.




Figure 3 | Current understanding of the mechanistic consequences of spliceosomal gene mutations for RNA splicing. **a** | Splicing factor 3B, subunit 1 (*SF3B1*) mutations (mutant form shown in red) alter 3' splice site (ss) selection by permitting or enhancing recognition of cryptic upstream 3' splice sites. It is not yet known how mutations affecting *SF3B1* alter its target protein–RNA and/or protein–protein interactions to drive this cryptic splice site recognition. Sequence motifs are illustrated as genomic DNA sequence rather than pre-mRNA sequence (T instead of U). **b** | U2 small nuclear RNA auxiliary factor 1 (*U2AF1*) mutations alter 3' ss consensus sequences. Wild-type *U2AF1* recognizes the consensus motif yAG|r at the intron–exon boundary (y = pyrimidine, r = purine, | = intron–exon boundary). S34F or S34Y mutations (shown in red) promote recognition of cAG|r over tAG|r, whereas Q157P or Q157R mutations (shown in red) promote recognition of yAG|g over yAG|a. **c** | Zinc finger, RNA-binding motif and serine/arginine-rich 2 (*ZRSR2*) mutations (shown in red) cause loss of *ZRSR2* function to induce splicing defects, primarily involving the aberrant retention of U12-type introns. **d** | Serine/arginine-rich splicing factor 2 (*SRSF2*) mutations (shown in red) alter exonic splicing enhancer preferences. Wild-type *SRSF2* recognizes the consensus motif SSNG (S = C or G), whereas mutant *SRSF2* preferentially recognizes the CCNG motif over the GGNG motif.

EZH2 occur in MDS^{99,100}, and *Ezh2* loss has been functionally linked to MDS development and aberrant haematopoietic stem cell self-renewal *in vivo*¹⁰¹. Therefore, decreased *EZH2* levels may partially explain how *SRSF2* mutations drive MDS, and may also explain the previously observed mutual exclusivity of *SRSF2* and *EZH2* mutations in MDS^{70,102}.

Mutations affecting other splicing factors. In addition to *SF3B1*, *SRSF2*, *U2AF1* and *ZRSR2*, other genes encoding splicing factors are recurrently mutated in cancer. Pre-mRNA processing factor 8 (*PRPF8*) is subjected to mutations or hemizygous deletions in 1–5% of

patients with myeloid leukaemias¹⁰³. Biochemical studies in yeast suggest that *PRPF8* mutations may affect recognition of suboptimal 3' splice sites¹⁰³.

Genes encoding splicing factors have also been identified as recurrent targets of translocations in cancer. For example, splicing factor proline/glutamine-rich (*SFPQ*; also known as *PSF*) is reportedly recurrently fused to *ABL1* in B-cell acute lymphoblastic leukaemia¹⁰⁴ and to transcription factor binding to IGHM enhancer 3 (*TFE3*) in papillary renal cell carcinomas¹⁰⁵. *SRSF3* is also reportedly involved in rare fusions with *BCL6* in B-cell lymphomas¹⁰⁶. Currently, it is not known whether these fusions affect the function of the splicing factors

involved in the chimeric protein product. In the case of *SFPQ* fusions, the sequence encoding the coiled-coil domain (which is important for protein dimerization) of *SFPQ* seems to be consistently included in the chimeric transcript¹⁰⁷, suggesting that *SFPQ* fusions may contribute to cancer by promoting aberrant dimerization of the partner protein of *SFPQ*.

Inherited genetic variation affecting RNA splicing factors has also been implicated in cancer. A missense genetic variant in serine/arginine repetitive matrix 2 (*SRRM2*; also known as *SRm300*) was recently found to segregate with incidence of familial papillary thyroid carcinoma¹⁰⁸. Patients carrying this *SRRM2* variant exhibited mis-splicing of specific cassette exons, suggesting that the variant altered the normal function of *SRRM2* in splicing. Somatic mutations may also interact with genetic variants to promote cancer. Recent work identified both somatic mutations and genetic variants affecting DEAD-box helicase 41 (*DDX41*) that are associated with high-penetrance familial MDS and AML¹⁰⁹. Although the normal molecular role of *DDX41* is incompletely understood, mass spectrometry data indicated that it interacts with core spliceosome components and that the likely loss-of-function *DDX41* mutations perturb these interactions. Therefore, *DDX41* may play a part in RNA splicing that is disrupted by MDS- and AML-associated mutations, although that hypothesis remains to be tested.

Connections to other cellular processes

Splicing factor dysregulation, including spliceosomal mutations, may directly or indirectly affect many cellular processes in addition to RNA splicing. These processes include the maintenance of genome integrity, epigenetic regulation, transcription, nuclear export and translation-dependent mRNA decay.

Depletion of *SRSF1* or *SRSF2* gives rise to DNA damage and genomic instability via the formation of RNA:DNA hybrids ('R loops')^{110,111}, which expose the unhybridized DNA strand to DNA damage (FIG. 4a). Splicing factors are also linked to the DNA damage response. For example, *BRCA1* is reportedly physically associated with *SF3B1* and *U2AF1* specifically in response to DNA damage¹¹². A recent study proposed that the spliceosome is an effector of ataxia telangiectasia mutated (*ATM*) signalling, wherein DNA lesions displace spliceosomes, resulting in R loop formation and *ATM* activation¹¹³.

Splicing is also closely linked to epigenetic regulation. Nucleosomes are preferentially positioned over exons rather than introns^{114,115}. *SF3B1* is preferentially associated with nucleosomes positioned over exons, facilitating recognition of these exons¹¹⁵. Histone H3 lysine 36 trimethylation (H3K36me3) is further enriched over exons^{116,117}, and modulation of H3K36me3 can influence splice site choice¹¹⁸. Conversely, modifying splice site recognition can influence H3K36me3 deposition^{119,120}. The literature linking splicing and epigenetics is reviewed in more detail in REFS 121,122.

Connections between splicing and epigenetics may contribute to the oncogenic activity of spliceosomal mutations. As described above, mutant *SRSF2*

prevents haematopoiesis in part by promoting a non-functional isoform of *EZH2*, resulting in global decreases in H3K27me3 levels⁷¹. Mutant *U2AF1* promotes a cancer-associated isoform of the histone variant macro-H2A.1 (REFS 75,87). Connections between *SF3B1* mutations and epigenetic dysregulation have not been identified, but are plausible given the known links between splice site recognition and chromatin described above. However, further studies are needed to determine whether potential epigenetic changes caused by *U2AF1* and/or *SF3B1* mutations are important for cancer initiation and progression.

Splicing factor dysregulation may also affect transcription independently of epigenetic regulation. For example, *SRSF2* regulates transcriptional elongation in a sequence-specific manner via the 7SK complex that governs RNA polymerase II pause release¹²³. Similarly, hnRNP A1, hnRNP A2 and hnRNP K are linked to cancer, and hnRNP A1 and hnRNP A2 have reported roles in transcription elongation whereas hnRNP K is linked to transcription termination^{124,125}. Many splicing factors co-transcriptionally associate with the C-terminal domain of RNA polymerase II, thereby coupling transcription, splicing, cleavage and polyadenylation (reviewed in REF. 126).

Finally, molecular changes induced by splicing factor dysregulation may not be confined to the nucleus. Many SR proteins facilitate nuclear export of both unspliced and spliced mRNAs^{127,128}. *SRSF1* shuttles between the nucleus and cytoplasm and promotes cap-dependent translation of bound mRNAs in a eukaryotic translation initiation factor 4E (eIF4E)-dependent manner^{129,130}. *SRSF1* activates the mTOR signalling pathway, which is required for *SRSF1*-mediated cell transformation^{39,41}. Splicing itself promotes translation via the deposition of multiprotein exon junction complexes (EJCs) near exon-exon junctions¹³¹.

NMD provides a concrete example of a cytoplasmic process that is probably affected by cancer-associated mutations affecting *SF3B1*, *SRSF2*, *U2AF1* and *ZRSR2*, even though those proteins localize to the nucleus. NMD is a translation-dependent RNA surveillance process that degrades mRNAs containing premature termination codons (reviewed in REFS 132,133) (FIG. 4b). Splicing and NMD are closely linked in human cells for several reasons. First, many NMD substrates are generated by alternative splicing, whereby premature termination codons are introduced through inclusion of alternatively spliced sequence containing an in-frame premature stop codon or exclusion of sequence, resulting in a frameshift. Second, stop codons are recognized as premature by the NMD machinery if they lie more than 50 nucleotides upstream of a splice junction^{132,133}. The 50 nucleotide threshold arises because the translating ribosome dislodges EJCs and prevents EJC-facilitated activation of NMD. However, if the ribosome stalls sufficiently upstream of an EJC, then NMD factors are recruited and activated^{132,133}. Third, specific splicing factors, including *SRSF1*, regulator of differentiation 1 (*ROD1*; also known as *PTBP3*) and *PTB* can enhance or repress NMD^{134–136}.

RNA polymerase II pause release

The process by which RNA polymerase II that is paused (not actively transcribing) after the initiation of transcription is released, enabling transcriptional elongation.

Frameshift

The disruption of an open reading frame by the insertion or deletion of nucleotide sequence whose length is not a multiple of three.




Figure 4 | Links between splicing factors and diverse biological processes and potential methods for therapeutic manipulation of splicing. **a** | Changes in the abundance, post-translational modifications and/or subcellular localization of splicing factors such as serine/arginine-rich splicing factor 1 (SRSF1) can cause DNA damage or influence the DNA damage response (DDR). Splicing factors have been linked to DNA damage and the DDR both directly (for example, insufficient levels of splicing factors can cause R loop formation and genomic instability) and indirectly (for example, through downstream changes in splicing). **b** | The steps involved in nonsense-mediated decay (NMD) are shown. The exon junction complex (EJC; grey) is deposited upstream of exon-exon junctions on the processed mRNA, and is displaced by the ribosome (blue) during the pioneer (first) round of translation. Ribosome stalling at a premature termination codon (PTC) >50 nt upstream of an EJC promotes interactions between release factors (purple) and up-frameshift suppressor 1 (UPF1; green), recruitment of other NMD components (yellow) and RNA degradation by endo- and exonucleases (beige). In contrast, if only a normal termination codon (TC) is present, then all EJCs are displaced by the ribosome during the pioneer round of translation and NMD is not triggered. Red stop signs indicate stop codons. **c** | Compounds and oligonucleotides that can disrupt or modulate normal splicing catalysis or alter splice site recognition through distinct pathways include drugs affecting U2 small nuclear ribonucleoprotein (snRNP) function, formation and/or interaction with pre-mRNA (panel i), drugs affecting post-translational modifications of serine/arginine-rich (SR) proteins and potentially other splicing factors (panel ii) and oligonucleotides to manipulate specific mRNA isoforms that may be important in tumour maintenance (panel iii). CLK, CDC2-like kinase; SF3B1, splicing factor 3B, subunit 1; SRPK, SR protein kinase; U2AF, U2 auxiliary factor.

Human cells express an abundance of mRNAs containing premature termination codons (one-third of all alternatively spliced isoforms by one estimate¹³⁷), including the *EZH2* poison exon that is promoted by *SRSF2* mutations⁷¹. A subset of poison exons are among the most evolutionarily conserved elements in the human genome^{138–140}. These poison exons enable splicing factors to post-transcriptionally downregulate

or upregulate expression of specific genes, including the genes encoding many splicing factors themselves, which may explain the considerable sequence conservation of many poison exons^{138–140}.

Interestingly, in the earliest report of splicing factor mutations, genes involved in NMD were upregulated following overexpression of mutant U2AF1 (REF. 8), suggesting a potential link between spliceosomal mutations

and overproduction of NMD substrates. However, such high levels of NMD substrates have not been observed in subsequent studies of mutations affecting U2AF1 or other splicing factors.

The recent discovery of recurrent mutations in *UPF1*, which encodes an RNA helicase that is central to NMD, in pancreatic adenocarcinoma provided a genetic link between NMD and cancer¹⁴¹. The observed mutations induced abnormal *UPF1* splicing and skipping of sequence encoding core domains, potentially resulting in partial or complete loss of *UPF1* function, although further work is required to determine how these mutations affect global RNA surveillance. Deficiencies in various NMD factors have been previously linked to disorders including intellectual disability¹⁴², thrombocytopenia with absent radii syndrome¹⁴³ and muscular dystrophy¹⁴⁴.

Implications for therapy

Given the crucial roles of specific alternatively spliced isoforms in cancer biology^{145,146}, as well as the potentially increased sensitivity of cancers to global perturbation of splicing efficiency relative to normal cells^{147,148}, pharmacological modulation of splicing may represent an important therapeutic strategy. Spliceosomal gene mutations that cause alteration or gain of function are mutually exclusive of one another and are always co-expressed with a wild-type allele, suggesting that cells bearing spliceosomal mutations may be unable to tolerate further perturbations in splicing, and could therefore be preferentially sensitive to pharmacological splicing inhibition. Several compounds and oligonucleotides have been identified that can disrupt or modulate normal splicing catalysis or alter splice site recognition through distinct pathways (FIG. 4c).

Several compounds with antitumour activity were identified through natural product screens before the discovery of spliceosomal gene mutations. Synthetic analogues with higher stability, solubility and activity were subsequently developed, including E7107, meayamycin, spliceostatin A and sudemycins (reviewed in REFS 149,150). Biochemical studies identified SF3B1 as the likely target of these drugs, consistent with the observation that mutations affecting the R1074 residue of SF3B1 confer resistance to pladienolide and E7107 (REFS 151,152). Unfortunately, two separate Phase I clinical trials of E7107 revealed an unexpected and unexplained side effect of visual disturbances in 5% of subjects^{153,154}. Further efforts are needed to determine whether this was an on- or off-target effect of U2 snRNP inhibition *in vivo*. In the meantime, several preclinical studies are evaluating the utility and safety of sudemycins^{155,156} for cancer therapy.

Although the origin of general antitumour activities of splicing inhibitors is unknown, two studies provided evidence that MYC expression renders cells sensitive to compounds that inhibit 3' splice site recognition^{147,148}. More recently, Lee *et al.*¹⁵⁷ reported that the splicing inhibitor E7107 reduced the leukaemic burden and prolonged survival of mice carrying oncogene-driven myeloid leukaemias if the leukaemias had *Srsf2*

mutations, but not if the leukaemias expressed only wild-type *Srsf2*. Lee *et al.*¹⁵⁷ observed similarly specific targeting of patient-derived xenograft (PDX) models of leukaemias with spliceosomal mutations. These data suggest that splicing inhibitors such as E7107 are synthetically lethal with genetic lesions affecting the spliceosome.

Interventions that target post-translational modification of splicing factors might also prove effective for therapy. For example, SR proteins are phosphorylated by kinases that include topoisomerase I and members of the SR protein kinase (SRPK) and CDC2-like kinase (CLK) families^{158–160}. These kinases affect SR protein subcellular localization and splicing activity^{161,162}, exhibit altered expression and/or activity in cancer^{163,164} and can potentially act as oncogenes¹⁶⁴. Small molecules that block activity of these kinases have been identified, including TG003, an inhibitor of CLK1 and CLK4, and SRPIN340, an SRPK1 and SRPK2 inhibitor that inhibits angiogenesis^{165,166}. In addition, indole derivatives, such as benzopyridoindoles and pyridocarbazoles, are a class of compounds that were recently discovered to modulate splicing by altering the ESE-dependent splicing activity of individual SR proteins¹⁶⁷. Indole derivatives have been shown to modulate the splicing event that generates the cancer-associated, constitutively active Δ RON isoform of the *RON* (also known as *MST1R*) proto-oncogene and revert the invasive phenotype of cancer cells expressing Δ RON¹⁶⁸.

Summary and future perspectives

The recent discovery of recurrent spliceosomal mutations as likely cancer drivers has underscored the pressing need to identify connections between abnormal pre-mRNA processing and tumorigenesis. Emerging evidence supports a model in which many spliceosomal mutations induce specific changes in splice site or exon recognition, frequently via altered RNA binding, leading to genome-wide splicing changes that presumably promote cancer development.

Despite these mechanistic advances, efforts to link altered splice site or exon recognition to specific pathological splicing events are nascent. Challenges include identification and prioritization among hundreds of downstream mis-spliced isoforms, as well as determination of the biological roles of specific isoforms. Furthermore, it is not known whether the pro-tumorigenic effects of mutated spliceosomal proteins are mediated by just a handful of mis-spliced isoforms, or instead are due to many splicing changes, which may even be functionally interdependent. Although many mis-spliced isoforms have been identified in cells bearing spliceosomal mutations, to date, very few of these isoforms have been functionally characterized.

Spliceosomal mutations probably both indirectly and directly dysregulate diverse cellular processes. In principle, spliceosomal mutations could affect almost any biological process by inducing mis-splicing of key regulators (for example, the connection between *SRSF2* and H3K27me3 deficiency via *EZH2* mis-splicing).

Spliceosomal mutations may also dysregulate processes such as transcriptional elongation, the DNA damage response and NMD, in which splicing factors have key roles (FIG. 4a,b).

Although spliceosomal mutations provide the most direct link between splicing and cancer, it is also important to note that abnormal splicing is a feature of most cancers even in the absence of spliceosomal mutations^{5–7}. Abnormal cancer-associated splicing may result from both specific and global perturbations to the splicing machinery. Specific perturbations may arise from dysregulation of single splicing factors that have a protumorigenic or antitumorigenic role (TABLE 1), whereas global perturbations may arise from effects including potential transcriptional amplification driven by MYC¹⁴⁸ or mutations affecting epigenetic regulators such as isocitrate dehydrogenase (IDH) or SET domain containing 2 (SETD2)^{5,7}.

Although incomplete, our current understanding of spliceosomal mutations suggests that these mutations may create new therapeutic opportunities. Because splicing factors can act as both oncoproteins and tumour suppressors, distinct therapeutic interventions may prove necessary for treating cancers harbouring different spliceosomal mutations. Possible therapeutic interventions fall into several broad categories, including restoring normal splicing and exploiting vulnerabilities to specifically target mutant cells.

Normal splicing could potentially be restored by specifically inhibiting the mutant protein, manipulating downstream splicing events or other methods. These approaches are promising, yet each requires further investigation. For example, specific inhibition or sequestration of mutant SRSF2 and U2AF1 may be possible given their altered RNA-binding preferences. However, definitive evidence that cancer cells depend on these mutated proteins, or that inhibiting the mutant allele is sufficient to restore normal splicing, is currently absent. Mutant SRSF2 and U2AF1 probably act as oncoproteins to promote tumour

formation, yet may not be required for subsequent tumour maintenance or growth. The same caveat applies to inhibition of downstream mis-splicing. Specific mis-splicing events could potentially be corrected with antisense oligonucleotides, which have shown promise in clinical trials of disorders such as Duchenne muscular dystrophy¹⁶⁹ and spinal muscular atrophy^{170,171}. However, our current understanding of how spliceosomal mutations perturb cellular function is insufficient to determine which mis-splicing events to correct in cancer. Furthermore, because inhibiting a mutant oncoprotein is likely to be more feasible than restoring the function of a disabled wild-type protein, restoring normal splicing may not be possible in the context of spliceosomal mutations that disable tumour suppressors. For example, ZRSR2 mutations cause loss of ZRSR2 expression or function, and it is unclear whether restoring U12-type intron recognition in the absence of ZRSR2 is possible.

Conversely, it may be feasible to selectively target cells expressing mutated splicing factors. Recent work suggested that inhibiting splicing catalysis itself may provide a therapeutic index in cells bearing spliceosomal mutations^{157,172}. Non-cell autonomous approaches to targeting cells with spliceosomal mutations may also be possible. Just as increased somatic mutational burdens may generate neoepitopes and render specific subsets of cancer sensitive to cancer immunotherapies^{173–176}, so may abnormal mRNAs generated by spliceosomal mutations result in neoepitope production in cancers bearing these lesions. Notably, these two approaches — inhibition of splicing catalysis and immunotherapy — could potentially be efficacious in the context of spliceosomal mutations that generate oncoproteins as well as those that inactivate tumour suppressors. Recurrent mutations in *SF3B1*, *SRSF2*, *U2AF1* and *ZRSR2* cause very different mechanistic alterations in splicing, yet each may render cells susceptible to further perturbation of splicing catalysis or result in the generation of neoepitopes.

- Zhang, J. & Manley, J. L. Misregulation of pre-mRNA alternative splicing in cancer. *Cancer Discov.* **3**, 1228–1237 (2013).
- David, C. J. & Manley, J. L. Alternative pre-mRNA splicing regulation in cancer: pathways and programs unhinged. *Genes Dev.* **24**, 2343–2364 (2010).
- Supek, F., Miñana, B., Valcárcel, J., Gabaldón, T. & Lehner, B. Synonymous mutations frequently act as driver mutations in human cancers. *Cell* **156**, 1324–1335 (2014).
- Jung, H. *et al.* Intron retention is a widespread mechanism of tumor-suppressor inactivation. *Nat. Genet.* **47**, 1242–1248 (2015).
- Dvinge, H. & Bradley, R. K. Widespread intron retention diversifies most cancer transcriptomes. *Genome Med.* **7**, 45 (2015).
- Danan-Gottthold, M. *et al.* Identification of recurrent regulated alternative splicing events across human solid tumors. *Nucleic Acids Res.* **43**, 5130–5144 (2015).
- Simon, J. M. *et al.* Variation in chromatin accessibility in human kidney cancer links H3K36 methyltransferase loss with widespread RNA processing defects. *Genome Res.* **24**, 241–250 (2014).
- Yoshida, K. *et al.* Frequent pathway mutations of splicing machinery in myelodysplasia. *Nature* **478**, 64–69 (2011).
- A landmark study demonstrating frequent mutations in genes encoding spliceosomal proteins in myeloid malignancies.**
- Graubert, T. A. *et al.* Recurrent mutations in the U2AF1 splicing factor in myelodysplastic syndromes. *Nat. Genet.* **44**, 53–57 (2012).
- Papaemmanuil, E. *et al.* Somatic *SF3B1* mutation in myelodysplasia with ring sideroblasts. *N. Engl. J. Med.* **365**, 1384–1395 (2011).
- Wang, L. *et al.* *SF3B1* and other novel cancer genes in chronic lymphocytic leukemia. *N. Engl. J. Med.* **365**, 2497–2506 (2011).
- A key study revealing frequent mutations in the gene encoding the spliceosomal protein SF3B1 in CLL.**
- Quesada, V. *et al.* Exome sequencing identifies recurrent mutations of the splicing factor *SF3B1* gene in chronic lymphocytic leukemia. *Nat. Genet.* **44**, 47–52 (2012).
- Pan, Q., Shai, O., Lee, L. J., Frey, B. J. & Blencowe, B. J. Deep surveying of alternative splicing complexity in the human transcriptome by high-throughput sequencing. *Nat. Genet.* **40**, 1413–1415 (2008).
- Wang, E. T. *et al.* Alternative isoform regulation in human tissue transcriptomes. *Nature* **456**, 470–476 (2008).
- A landmark study describing the use of RNA-seq to quantify splicing across human tissues.**
- Wahl, M. C., Will, C. L. & Lührmann, R. The spliceosome: design principles of a dynamic RNP machine. *Cell* **136**, 701–718 (2009).
- Turunen, J. J., Niemelä, E. H., Verma, B. & Frilander, M. J. The significant other: splicing by the minor spliceosome. *Wiley Interdiscip. Rev. RNA* **4**, 61–76 (2013).
- Matera, A. G. & Wang, Z. A day in the life of the spliceosome. *Nat. Rev. Mol. Cell Biol.* **15**, 108–121 (2014).
- Fica, S. M. *et al.* RNA catalyses nuclear pre-mRNA splicing. *Nature* **503**, 229–234 (2013).
- Graveley, B. R., Hertel, K. J. & Maniatis, T. A systematic analysis of the factors that determine the strength of pre-mRNA splicing enhancers. *EMBO J.* **17**, 6747–6756 (1998).
- Wang, Z. & Burge, C. B. Splicing regulation: from a parts list of regulatory elements to an integrated splicing code. *RNA* **14**, 802–813 (2008).
- Fu, X. D. & Ares, M. Jr. Context-dependent control of alternative splicing by RNA-binding proteins. *Nat. Rev. Genet.* **15**, 689–701 (2014).
- Singh, R. & Valcárcel, J. Building specificity with nonspecific RNA-binding proteins. *Nat. Struct. Mol. Biol.* **12**, 645–653 (2005).

23. Long, J. C. & Caceres, J. F. The SR protein family of splicing factors: master regulators of gene expression. *Biochem. J.* **417**, 15–27 (2009).
24. Zhou, Z. & Fu, X. D. Regulation of splicing by SR proteins and SR protein-specific kinases. *Chromosoma* **122**, 191–207 (2013).
25. Krecic, A. M. & Swanson, M. S. hnRNP complexes: composition, structure, and function. *Curr. Opin. Cell Biol.* **11**, 363–371 (1999).
26. Zahler, A. M., Lane, W. S., Stolk, J. A. & Roth, M. B. SR proteins: a conserved family of pre-mRNA splicing factors. *Genes Dev.* **6**, 837–847 (1992).
27. Kohtz, J. D. *et al.* Protein–protein interactions and 5′-splice-site recognition in mammalian mRNA precursors. *Nature* **368**, 119–124 (1994).
28. Pandit, S. *et al.* Genome-wide analysis reveals SR protein cooperation and competition in regulated splicing. *Mol. Cell* **50**, 223–235 (2013).
29. Han, J. *et al.* SR proteins induce alternative exon skipping through their activities on the flanking constitutive exons. *Mol. Cell Biol.* **31**, 793–802 (2011).
30. Xue, Y. *et al.* Genome-wide analysis of PTB–RNA interactions reveals a strategy used by the general splicing repressor to modulate exon inclusion or skipping. *Mol. Cell* **36**, 996–1006 (2009).
31. Llorian, M. *et al.* Position-dependent alternative splicing activity revealed by global profiling of alternative splicing events regulated by PTB. *Nat. Struct. Mol. Biol.* **17**, 1114–1123 (2010).
32. Huelga, S. C. *et al.* Integrative genome-wide analysis reveals cooperative regulation of alternative splicing by hnRNP proteins. *Cell Rep.* **1**, 167–178 (2012).
33. Dasgupta, T. & Ladd, A. N. The importance of CELF control: molecular and biological roles of the CUG-BP, Elav-like family of RNA-binding proteins. *Wiley Interdiscip. Rev. RNA* **3**, 104–121 (2012).
34. Konieczny, P., Stepniak-Konieczna, E. & Sobczak, K. MBNL proteins and their target RNAs, interaction and splicing regulation. *Nucleic Acids Res.* **42**, 10873–10887 (2014).
35. Ule, J. *et al.* An RNA map predicting Nova-dependent splicing regulation. *Nature* **444**, 580–586 (2006).
36. Saltzman, A. L., Pan, Q. & Blencowe, B. J. Regulation of alternative splicing by the core spliceosomal machinery. *Genes Dev.* **25**, 373–384 (2011).
37. Jurica, M. S. & Moore, M. J. Pre-mRNA splicing: awash in a sea of proteins. *Mol. Cell* **12**, 5–14 (2003).
38. Barash, Y. *et al.* Deciphering the splicing code. *Nature* **465**, 53–59 (2010).
39. Karni, R. *et al.* The gene encoding the splicing factor SF2/ASF is a proto-oncogene. *Nat. Struct. Mol. Biol.* **14**, 185–193 (2007).
- This paper demonstrated that modest overexpression of the splicing factor SRSF1 is pro-tumorigenic.**
40. Anczuków, O. *et al.* The splicing factor SRSF1 regulates apoptosis and proliferation to promote mammary epithelial cell transformation. *Nat. Struct. Mol. Biol.* **19**, 220–228 (2012).
41. Karni, R., Hippo, Y., Lowe, S. W. & Krainer, A. R. The splicing-factor oncoprotein SF2/ASF activates mTORC1. *Proc. Natl Acad. Sci. USA* **105**, 15323–15327 (2008).
42. Jia, R., Li, C., McCoy, J. P., Deng, C.-X. X. & Zheng, Z.-M. M. SRp20 is a proto-oncogene critical for cell proliferation and tumor induction and maintenance. *Int. J. Biol. Sci.* **6**, 806–826 (2010).
43. Tang, Y. *et al.* Downregulation of splicing factor SRSF3 induces p53 β , an alternatively spliced isoform of p53 that promotes cellular senescence. *Oncogene* **32**, 2792–2798 (2013).
44. Jensen, M. A., Wilkinson, J. E. & Krainer, A. R. Splicing factor SRSF6 promotes hyperplasia of sensitized skin. *Nat. Struct. Mol. Biol.* **21**, 189–197 (2014).
45. Cohen-Elav, M. *et al.* The splicing factor SRSF6 is amplified and is an oncoprotein in lung and colon cancers. *J. Pathol.* **229**, 630–639 (2013).
46. David, C. J., Chen, M., Assanah, M., Canoll, P. & Manley, J. L. hnRNP proteins controlled by c-Myc deregulate pyruvate kinase mRNA splicing in cancer. *Nature* **463**, 364–368 (2010).
47. Clower, C. V. *et al.* The alternative splicing repressors hnRNP A1/A2 and PTB influence pyruvate kinase isoform expression and cell metabolism. *Proc. Natl Acad. Sci. USA* **107**, 1894–1899 (2010).
48. Babic, I. *et al.* EGFR mutation-induced alternative splicing of Max contributes to growth of glycolytic tumors in brain cancer. *Cell Metab.* **17**, 1000–1008 (2013).
49. Golan-Gerstl, R. *et al.* Splicing factor hnRNP A2/B1 regulates tumor suppressor gene splicing and is an oncogenic driver in glioblastoma. *Cancer Res.* **71**, 4464–4472 (2011).
50. Sweetser, D. A. *et al.* Delineation of the minimal commonly deleted segment and identification of candidate tumor-suppressor genes in del(9q) acute myeloid leukemia. *Genes Chromosomes Cancer* **44**, 279–291 (2005).
51. Gallardo, M. *et al.* hnRNP K is a haploinsufficient tumor suppressor that regulates proliferation and differentiation programs in hematologic malignancies. *Cancer Cell* **28**, 486–499 (2015).
52. Moumen, A., Masterson, P., O'Connor, M. J. & Jackson, S. P. hnRNP K: an HDM2 target and transcriptional coactivator of p53 in response to DNA damage. *Cell* **123**, 1065–1078 (2005).
53. Zong, F. Y. *et al.* The RNA-binding protein QKI suppresses cancer-associated aberrant splicing. *PLoS Genet.* **10**, e1004289 (2014).
54. Angeloni, D. Molecular analysis of deletions in human chromosome 3p21 and the role of resident cancer genes in disease. *Brief. Funct. Genomic. Proteomic.* **6**, 19–39 (2007).
55. Imielinski, M. *et al.* Mapping the hallmarks of lung adenocarcinoma with massively parallel sequencing. *Cell* **150**, 1107–1120 (2012).
56. Oh, J. J. *et al.* 3p21.3 tumor suppressor gene H37/Luca15/RBM5 inhibits growth of human lung cancer cells through cell cycle arrest and apoptosis. *Cancer Res.* **66**, 3419–3427 (2006).
57. Rintala-Maki, N. D. & Goard, C. A. Expression of RBM5-related factors in primary breast tissue. *J. Cell. Biochem.* **100**, 1440–1458 (2007).
58. Hernandez, J. *et al.* Tumor suppressor properties of the splicing regulatory factor RBM10. *RNA Biol.* **13**, 466–472 (2016).
59. Bechara, E. G., Sebestyén, E., Bernardis, I., Eyra, E. & Valcárcel, J. RBM5, 6, and 10 differentially regulate NUMB alternative splicing to control cancer cell proliferation. *Mol. Cell* **52**, 720–733 (2013).
60. Wang, Y. *et al.* The splicing factor RBM4 controls apoptosis, proliferation, and migration to suppress tumor progression. *Cancer Cell* **26**, 374–389 (2014).
61. Warzecha, C. C., Sato, T. K., Nabet, B., Hogenesch, J. B. & Carstens, R. P. ESRP1 and ESRP2 are epithelial cell-type-specific regulators of FGFR2 splicing. *Mol. Cell* **35**, 591–601 (2009).
62. Shapiro, I. M. *et al.* An EMT-driven alternative splicing program occurs in human breast cancer and modulates cellular phenotype. *PLoS Genet.* **7**, e1002218 (2011).
63. Harbour, J. W. *et al.* Recurrent mutations at codon 625 of the splicing factor SF3B1 in uveal melanoma. *Nat. Genet.* **45**, 133–135 (2013).
64. Martin, M. *et al.* Exome sequencing identifies recurrent somatic mutations in *ELF1A* and *SF3B1* in uveal melanoma with disomy 3. *Nat. Genet.* **45**, 933–936 (2013).
65. Biankin, A. V. *et al.* Pancreatic cancer genomes reveal aberrations in axon guidance pathway genes. *Nature* **491**, 399–405 (2012).
66. Stephens, P. J. *et al.* The landscape of cancer genes and mutational processes in breast cancer. *Nature* **486**, 400–404 (2012).
67. Ellis, M. J. *et al.* Whole-genome analysis informs breast cancer response to aromatase inhibition. *Nature* **486**, 353–360 (2012).
68. Maguire, S. L. *et al.* SF3B1 mutations constitute a novel therapeutic target in breast cancer. *J. Pathol.* **235**, 571–580 (2015).
69. Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. *Nature* **490**, 61–70 (2012).
70. Papaemmanuil, E. *et al.* Clinical and biological implications of driver mutations in myelodysplastic syndromes. *Blood* **122**, 3616–3627 (2013).
71. Kim, E. *et al.* SRSF2 mutations contribute to myelodysplasia by mutant-specific effects on exon recognition. *Cancer Cell* **27**, 617–630 (2015).
- This study demonstrated that SRSF2 mutations are sufficient to drive myelodysplasia, are distinct from loss-of-function of SRSF2 and change the RNA-binding affinity of the protein to alter exon recognition.**
72. Kon, A. *et al.* SRSF2 P95H mutation causes impaired stem cell repopulation and hematopoietic differentiation in mice. *Blood Abstr.* **126**, 1649 (2015).
73. Obeng, E. A. *et al.* Mutant splicing factor 3b subunit 1 (SF3B1) causes dysregulated erythropoiesis and a stem cell disadvantage. *Blood Abstr.* **124**, 828 (2014).
74. Mupo, A. *et al.* SF3b1 K700E mutation impairs pre-mRNA splicing and definitive hematopoiesis in a conditional knock-in mouse model. *Blood Abstr.* **126**, 140 (2015).
75. Shirai, C. L. *et al.* Mutant U2AF1 expression alters hematopoiesis and pre-mRNA splicing *in vivo*. *Cancer Cell* **27**, 631–643 (2015).
- This study presented one of the first *in vivo* models of spliceosomal gene mutations in cancer, which revealed the biological effects of the U2AF1^{S54F} mutation on hematopoiesis.**
76. Mian, S. A. *et al.* SF3B1 mutant MDS-initiating cells may arise from the hematopoietic stem cell compartment. *Nat. Commun.* **6**, 10004 (2015).
77. Lindsley, R. C. *et al.* Acute myeloid leukemia ontogeny is defined by distinct somatic mutations. *Blood* **125**, 1367–1376 (2015).
78. Rossi, D. *et al.* Mutations of the SF3B1 splicing factor in chronic lymphocytic leukemia: association with progression and fludarabine-refractoriness. *Blood* **118**, 6904–6908 (2011).
79. DeBoever, C. *et al.* Transcriptome sequencing reveals potential mechanism of cryptic 3′ splice site selection in SF3B1-mutated cancers. *PLoS Comput. Biol.* **11**, e1004105 (2015).
80. Darman, R. B. *et al.* Cancer-associated SF3B1 hotspot mutations induce cryptic 3′ splice site selection through use of a different branch point. *Cell Rep.* **13**, 1033–1045 (2015).
- This study revealed that SF3B1 mutations promote recognition of cryptic 3′ splice sites in cancer samples and genetically modified cell lines.**
81. Allikmets, R. *et al.* Mutation of a putative mitochondrial iron transporter gene (*ABC7*) in X-linked sideroblastic anemia and ataxia (XLSA/A). *Hum. Mol. Genet.* **8**, 743–749 (1999).
82. Ponder, C. *et al.* *Abcb7*, the gene responsible for X-linked sideroblastic anemia with ataxia, is essential for hematopoiesis. *Blood* **109**, 3567–3569 (2007).
83. Wu, S., Romfo, C. M., Nilsen, T. W. & Green, M. R. Functional recognition of the 3′ splice site AG by the splicing factor U2AF35. *Nature* **402**, 832–835 (1999).
84. Reed, R. The organization of 3′ splice-site sequences in mammalian introns. *Genes Dev.* **3**, 2113–2123 (1989).
85. Przychodzen, B. *et al.* Patterns of missplicing due to somatic U2AF1 mutations in myeloid neoplasms. *Blood* **122**, 999–1006 (2013).
86. Brooks, A. N. *et al.* A pan-cancer analysis of transcriptome changes associated with somatic mutations in U2AF1 reveals commonly altered splicing events. *PLoS ONE* **9**, e87361 (2014).
87. Ilagan, J. O. *et al.* U2AF1 mutations alter splice site recognition in hematological malignancies. *Genome Res.* **25**, 14–26 (2015).
- This study demonstrated that U2AF1 mutations cause alteration, not loss, of function and that mutations in the first versus second zinc fingers of U2AF1 induce different changes in 3′ splice site preference.**
88. Okey-Owuor, T. *et al.* U2AF1 mutations alter sequence specificity of pre-mRNA binding and splicing. *Leukemia* **29**, 909–917 (2015).
89. Damm, F. *et al.* *BCOR* and *BCORL1* mutations in myelodysplastic syndromes and related disorders. *Blood* **122**, 3169–3177 (2013).
90. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature* **517**, 576–582 (2015).
91. Shen, H., Zheng, X., Luecke, S. & Green, M. R. The U2AF35-related protein Urp contacts the 3′ splice site to promote U12-type intron splicing and the second step of U2-type intron splicing. *Genes Dev.* **24**, 2389–2394 (2010).
92. Madan, V. *et al.* Aberrant splicing of U12-type introns is the hallmark of ZRSR2 mutant myelodysplastic syndrome. *Nat. Commun.* **6**, 6042 (2015).
- This work demonstrated that U12-type introns are poorly recognized in patients carrying presumed loss-of-function ZRSR2 mutations.**
93. Bejar, R. *et al.* Validation of a prognostic model and the impact of mutations in patients with lower-risk myelodysplastic syndromes. *J. Clin. Oncol.* **30**, 3376–3382 (2012).
94. Graveley, B. R. & Maniatis, T. Arginine/serine-rich domains of SR proteins can function as activators of pre-mRNA splicing. *Mol. Cell* **1**, 765–771 (1998).
95. Liu, H. X., Chew, S. L., Cartegni, L., Zhang, M. Q. & Krainer, A. R. Exonic splicing enhancer motif recognized by human SC35 under splicing conditions. *Mol. Cell Biol.* **20**, 1063–1071 (2000).

96. Schaaf, T. D. & Maniatis, T. Multiple distinct splicing enhancers in the protein-coding sequences of a constitutively spliced pre-mRNA. *Mol. Cell. Biol.* **19**, 261–273 (1999).
97. Zhang, J. *et al.* Disease-associated mutation in *SRSF2* misregulates splicing by altering RNA-binding affinities. *Proc. Natl Acad. Sci. USA* **112**, E4726–E4734 (2015).
98. Daubner, G. M., Clery, A., Jayne, S., Stevenin, J. & Allain, F. H. A syn-anti conformational difference allows SRSF2 to recognize guanines and cytosines equally well. *EMBO J.* **31**, 162–174 (2012).
- This study described NMR solution structures of the RRM domain of SRSF2 in complex with RNA that explained the ability of SRSF2 to bind to both C-rich and C-rich motifs.**
99. Ernst, T. *et al.* Inactivating mutations of the histone methyltransferase gene *EZH2* in myeloid disorders. *Nat. Genet.* **42**, 722–726 (2010).
100. Nikolski, G. *et al.* Somatic mutations of the histone methyltransferase gene *EZH2* in myelodysplastic syndromes. *Nat. Genet.* **42**, 665–667 (2010).
101. Muto, T. *et al.* Concurrent loss of *Ezh2* and *Tet2* cooperates in the pathogenesis of myelodysplastic disorders. *J. Exp. Med.* **210**, 2627–2639 (2013).
102. Haeflrich, T. *et al.* Landscape of genetic lesions in 944 patients with myelodysplastic syndromes. *Leukemia* **28**, 241–247 (2014).
103. Kurtovic-Kozaric, A. *et al.* PRPF8 defects cause missplicing in myeloid malignancies. *Leukemia* **29**, 126–136 (2015).
104. Duhoux, F. P. *et al.* The t(1;9)(p34;q34) fusing *ABL1* with *SFPQ*, a pre-mRNA processing gene, is recurrent in acute lymphoblastic leukemias. *Leukemia Res.* **35**, e114–e117 (2011).
105. Mathur, M. & Samuels, H. H. Role of PSF-TFE3 oncoprotein in the development of papillary renal cell carcinomas. *Oncogene* **26**, 277–283 (2007).
106. Chen, W., Itoyama, T. & Chaganti, R. S. Splicing factor SRP20 is a novel partner of BCL6 in a t(3;6)(q27;p21) translocation in transformed follicular lymphoma. *Genes Chromosomes Cancer* **32**, 281–284 (2001).
107. Lee, M. *et al.* The structure of human SFPQ reveals a coiled-coil mediated polymer essential for functional aggregation in gene regulation. *Nucleic Acids Res.* **43**, 3826–3840 (2015).
108. Tomasic, J. *et al.* A germline mutation in *SRRM2*, a splicing factor gene, is implicated in papillary thyroid carcinoma predisposition. *Sci. Rep.* **5**, 10566 (2015).
109. Polprasert, C. *et al.* Inherited and somatic defects in *DDX41* in myeloid neoplasms. *Cancer Cell* **27**, 658–670 (2015).
110. Li, X. & Manley, J. L. Inactivation of the SR protein splicing factor ASF/SF2 results in genomic instability. *Cell* **122**, 365–378 (2005).
- This study demonstrated that loss of SRSF1 results in R loop formation and DNA damage.**
111. Xiao, R. *et al.* Splicing regulator SC35 is essential for genomic stability and cell proliferation during mammalian organogenesis. *Mol. Cell. Biol.* **27**, 5393–5402 (2007).
112. Savage, K. I. *et al.* Identification of a BRCA1-mRNA splicing complex required for efficient DNA repair and maintenance of genomic stability. *Mol. Cell* **54**, 445–459 (2014).
113. Tresini, M. *et al.* The core spliceosome as target and effector of non-canonical ATM signalling. *Nature* **523**, 53–58 (2015).
114. Tilgner, H. *et al.* Nucleosome positioning as a determinant of exon recognition. *Nat. Struct. Mol. Biol.* **16**, 996–1001 (2009).
115. Kfir, N. *et al.* SF3B1 association with chromatin determines splicing outcomes. *Cell Rep.* **11**, 618–629 (2015).
116. Kolasinska-Zwiercz, P. *et al.* Differential chromatin marking of introns and expressed exons by H3K36me3. *Nat. Genet.* **41**, 376–381 (2009).
117. Spies, N., Nielsen, C. B., Padgett, R. A. & Burge, C. B. Biased chromatin signatures around polyadenylation sites and exons. *Mol. Cell* **36**, 245–254 (2009).
118. Luco, R. F. *et al.* Regulation of alternative splicing by histone modifications. *Science* **327**, 996–1000 (2010).
- This paper demonstrated that histone modifications can influence alternative splicing.**
119. De Almeida, S. F. F. *et al.* Splicing enhances recruitment of methyltransferase HYPB/Setd2 and methylation of histone H3 Lys36. *Nat. Struct. Mol. Biol.* **18**, 977–983 (2011).
120. Kim, S., Kim, H., Fong, N., Erickson, B. & Bentley, D. L. Pre-mRNA splicing is a determinant of histone H3K36 methylation. *Proc. Natl Acad. Sci. USA* **108**, 13564–13569 (2011).
121. Luco, R. F., Allo, M., Schor, I. E., Kornblihtt, A. R. & Misteli, T. Epigenetics in alternative pre-mRNA splicing. *Cell* **144**, 16–26 (2011).
122. Khan, D. H., Jahan, S. & Davie, J. R. Pre-mRNA splicing: role of epigenetics and implications in disease. *Adv. Biol. Regul.* **52**, 377–388 (2012).
123. Ji, X. *et al.* SR proteins collaborate with 7SK and promoter-associated nascent RNA to release paused polymerase. *Cell* **153**, 855–868 (2013).
124. Lemieux, B. *et al.* A function for the hnRNP A1/A2 proteins in transcription elongation. *PLoS ONE* **10**, e0126654 (2015).
125. Mikula, M., Bomsztyk, K., Coryca, K., Chojnowski, K. & Ostrowski, J. Heterogeneous nuclear ribonucleoprotein (hnRNP) K genome-wide binding survey reveals its role in regulating 3'-end RNA processing and transcription termination at the early growth response 1 (*EGR1*) gene through XRN2 exonuclease. *J. Biol. Chem.* **288**, 24788–24798 (2013).
126. Hsin, J. P. & Manley, J. L. The RNA polymerase II CTD coordinates transcription and RNA processing. *Genes Dev.* **26**, 2119–2137 (2012).
127. Huang, Y., Gattoni, R., Stevenin, J. & Steitz, J. A. SR splicing factors serve as adapter proteins for TAP-dependent mRNA export. *Mol. Cell* **11**, 837–843 (2003).
128. Muller-McNicoll, M. *et al.* SR proteins are NXF1 adaptors that link alternative RNA processing to mRNA export. *Genes Dev.* **30**, 553–566 (2016).
129. Sanford, J. R., Gray, N. K., Beckmann, K. & Caceres, J. F. A novel role for shuttling SR proteins in mRNA translation. *Genes Dev.* **18**, 755–768 (2004).
130. Michlewski, G., Sanford, J. R. & Caceres, J. F. The splicing factor SF2/ASF regulates translation initiation by enhancing phosphorylation of 4E-BP1. *Mol. Cell* **30**, 179–189 (2008).
131. Nott, A., Le Hir, H. & Moore, M. J. Splicing enhances translation in mammalian cells: an additional function of the exon junction complex. *Genes Dev.* **18**, 210–222 (2004).
132. Chang, Y.-F. F., Imam, J. S. & Wilkinson, M. F. The nonsense-mediated decay RNA surveillance pathway. *Annu. Rev. Biochem.* **76**, 51–74 (2007).
133. Popp, M. W. & Maquat, L. E. Organizing principles of mammalian nonsense-mediated mRNA decay. *Annu. Rev. Genet.* **47**, 139–165 (2013).
134. Zhang, Z. & Krainer, A. R. Involvement of SR proteins in mRNA surveillance. *Mol. Cell* **16**, 597–607 (2004).
135. Brazao, T. F. *et al.* A new function of ROD1 in nonsense-mediated mRNA decay. *FEBS Lett.* **586**, 1101–1110 (2012).
136. Ge, Z., Quek, B. L., Beemon, K. L. & Hogg, J. R. Polypyrimidine tract binding protein 1 protects mRNAs from recognition by the nonsense-mediated mRNA decay pathway. *eLife* **5**, e11155 (2016).
137. Lewis, B. P., Green, R. E. & Brenner, S. E. Evidence for the widespread coupling of alternative splicing and nonsense-mediated mRNA decay in humans. *Proc. Natl Acad. Sci. USA* **100**, 189–192 (2003).
- This study demonstrated that approximately one-third of alternative isoforms of human genes contain premature termination codons that probably trigger NMD.**
138. Lareau, L. F., Inada, M., Green, R. E., Wengrod, J. C. & Brenner, S. E. Unproductive splicing of SR genes associated with highly conserved and ultraconserved DNA elements. *Nature* **446**, 926–929 (2007).
139. Ni, J. Z. *et al.* Ultraconserved elements are associated with homeostatic control of splicing regulators by alternative splicing and nonsense-mediated decay. *Genes Dev.* **21**, 708–718 (2007).
140. Lareau, L. F. & Brenner, S. E. Regulation of splicing factors by alternative splicing and NMD is conserved between kingdoms yet evolutionarily flexible. *Mol. Biol. Evol.* **32**, 1072–1079 (2015).
141. Liu, C. *et al.* The *UPF1* RNA surveillance gene is commonly mutated in pancreatic adenocarcinoma. *Nat. Med.* **20**, 596–598 (2014).
142. Tarpey, P. S. *et al.* Mutations in *UPF3B*, a member of the nonsense-mediated mRNA decay complex, cause syndromic and nonsyndromic mental retardation. *Nat. Genet.* **39**, 1127–1133 (2007).
143. Albers, C. A. *et al.* Compound inheritance of a low-frequency regulatory SNP and a rare null mutation in exon-junction complex subunit RBM8A causes TAR syndrome. *Nat. Genet.* **44**, 435–439 (2012).
144. Feng, Q. *et al.* A feedback loop between nonsense-mediated decay and the retrograde DUX4 in facioscapulohumeral muscular dystrophy. *eLife* **4**, e04996 (2015).
145. Harper, S. J. & Bates, D. O. VEGF-A splicing: the key to anti-angiogenic therapeutics? *Nat. Rev. Cancer* **8**, 880–887 (2008).
146. Christoff, H. R. *et al.* The M2 splice isoform of pyruvate kinase is important for cancer metabolism and tumour growth. *Nature* **452**, 230–233 (2008).
147. Hubert, C. G. *et al.* Genome-wide RNAi screens in human brain tumor isolates reveal a novel viability requirement for PHF5A. *Genes Dev.* **27**, 1032–1045 (2013).
148. Hsu, T. Y. *et al.* The spliceosome is a therapeutic vulnerability in MYC-driven cancer. *Nature* **525**, 384–388 (2015).
149. Bonnal, S., Végivani, L. & Válcárcel, J. The spliceosome as a target of novel antitumour drugs. *Nat. Rev. Drug Discov.* **11**, 847–859 (2012).
150. Webb, T. R., Joyner, A. S. & Potter, P. M. The development and application of small molecule modulators of SF3b as therapeutic agents for cancer. *Drug Discov. Today* **18**, 43–49 (2013).
151. Yokoi, A. *et al.* Biological validation that SF3b is a target of the antitumor macrolide pladienolide. *FEBS J.* **278**, 4870–4880 (2011).
152. Kotake, Y., Sagane, K., Owa, T. & Mimori-Kiyosue, Y. Splicing factor SF3b as a target of the antitumor natural product pladienolide. *Nat. Chem. Biol.* **3**, 570–575 (2007).
153. Eskens, F. A. *et al.* Phase I pharmacokinetic and pharmacodynamic study of the first-in-class spliceosome inhibitor E7107 in patients with advanced solid tumors. *Clin. Cancer Res.* **19**, 6296–6304 (2013).
154. Hong, D. S. *et al.* A phase I, open-label, single-arm, dose-escalation study of E7107, a precursor messenger ribonucleic acid (pre-mRNA) spliceosome inhibitor administered intravenously on days 1 and 8 every 21 days to patients with solid tumors. *Invest. New Drugs* **32**, 436–444 (2013).
155. Fan, L., Lagisetty, C., Edwards, C. C., Webb, T. R. & Potter, P. M. Sudemycin, novel small molecule analogues of FR901464, induce alternative gene splicing. *ACS Chem. Biol.* **6**, 582–589 (2011).
156. Xargay-Torrent, S. *et al.* The splicing modulator sudemycin induces a specific antitumor response and cooperates with ibrutinib in chronic lymphocytic leukemia. *Oncotarget* **6**, 22734–22749 (2015).
157. Lee, S. C.-W. *et al.* Modulation of splicing catalysis for therapeutic targeting of leukemia with mutations in genes encoding spliceosomal proteins. *Nat. Med.* <http://dx.doi.org/10.1038/nm.4097> (2016).
158. Rossi, F. *et al.* Specific phosphorylation of SR proteins by mammalian DNA topoisomerase I. *Nature* **381**, 80–82 (1996).
159. Gui, J. F., Lane, W. S. & Fu, X. D. A serine kinase regulates intracellular localization of splicing factors in the cell cycle. *Nature* **369**, 678–682 (1994).
160. Colwill, K. *et al.* The Cln/Sty protein kinase phosphorylates SR splicing factors and regulates their intranuclear distribution. *EMBO J.* **15**, 265–275 (1996).
161. Stamm, S. Regulation of alternative splicing by reversible protein phosphorylation. *J. Biol. Chem.* **283**, 1223–1227 (2008).
162. Yeakley, J. M. *et al.* Phosphorylation regulates *in vivo* interaction and molecular targeting of serine/arginine-rich pre-mRNA splicing factors. *J. Cell Biol.* **145**, 447–455 (1999).
163. Gout, S. *et al.* Abnormal expression of the pre-mRNA splicing regulators SRSF1, SRSF2, SRPK1 and SRPK2 in non small cell lung carcinoma. *PLoS ONE* **7**, e46539 (2012).
164. Yoshida, T. *et al.* CLK2 is an oncogenic kinase and splicing regulator in breast cancer. *Cancer Res.* **75**, 1516–1526 (2015).
165. Dawid, G. N. *et al.* Regulation of vascular endothelial growth factor (VEGF) splicing from pro-angiogenic to anti-angiogenic isoforms a novel therapeutic strategy for angiogenesis. *J. Biol. Chem.* **19**, 5532–5540 (2010).
166. Elianna, M. A. *et al.* WT1 mutants reveal SRPK1 to be a downstream angiogenesis target by altering VEGF splicing. *Cancer Cell* **20**, 768–780 (2010).
167. Soret, J. *et al.* Selective modification of alternative splicing by indole derivatives that target serine-arginine-rich protein splicing factors. *Proc. Natl Acad. Sci. USA* **102**, 8764–8769 (2005).

168. Ghigna, C. *et al.* Pro-metastatic splicing of Ron proto-oncogene mRNA can be reversed: Therapeutic potential of bifunctional oligonucleotides and indole derivatives. *RNA Biol.* **7**, 495–503 (2010).
169. Aartsma-Rus, A., Fokkema, I. & Verschuuren, J. Theoretic applicability of antisense-mediated exon skipping for Duchenne muscular dystrophy mutations. *Hum. Mutat.* **30**, 293–299 (2009).
170. Hua, Y., Sahashi, K., Hung, G. & Rigo, F. Antisense correction of SMN2 splicing in the CNS rescues necrosis in a type III SMA mouse model. *Genes Dev.* **24**, 1634–1644 (2010).
171. Passini, M. A. *et al.* Antisense oligonucleotides delivered to the mouse CNS ameliorate symptoms of severe spinal muscular atrophy. *Sci. Transl. Med.* **3**, 72ra18 (2011).
172. Zhou, Q. *et al.* A chemical genetics approach for the functional assessment of novel cancer genes. *Cancer Res.* **75**, 1949–1958 (2015).
173. Snyder, A. *et al.* Genetic basis for clinical response to CTLA-4 blockade in melanoma. *N. Engl. J. Med.* **371**, 2189–2199 (2014).
174. Rizvi, N. A. *et al.* Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science* **348**, 124–128 (2015).
175. Le, D. T. *et al.* PD-1 blockade in tumors with mismatch-repair deficiency. *N. Engl. J. Med.* **372**, 2509–2520 (2015).
176. Gubin, M. M. *et al.* Checkpoint blockade cancer immunotherapy targets tumour-specific mutant antigens. *Nature* **515**, 577–581 (2014).
177. Alamancos, G. P., Agirre, E. & Eyra, E. Methods to study splicing from high-throughput RNA sequencing data. *Methods Mol. Biol.* **1126**, 357–397 (2014).
178. Engstrom, P. G. *et al.* Systematic evaluation of spliced alignment programs for RNA-seq data. *Nat. Methods* **10**, 1185–1191 (2013).
179. Thierry-Mieg, D. & Thierry-Mieg, J. AceView: a comprehensive cDNA-supported gene and transcripts annotation. *Genome Biol.* **7**, S12.1–S12.14 (2006).
180. Cunningham, F. *et al.* Ensembl 2015. *Nucleic Acids Res.* **43**, 9 (2015).
181. Rosenbloom, K. R. *et al.* The UCSC Genome Browser database: 2015 update. *Nucleic Acids Res.* **43**, 81 (2015).
182. Katz, Y., Wang, E. T., Airolidi, E. M. & Burge, C. B. Analysis and design of RNA sequencing experiments for identifying isoform regulation. *Nat. Methods* **7**, 1009–1015 (2010).
183. Barbosa-Morais, N. L. *et al.* The evolutionary landscape of alternative splicing in vertebrate species. *Science* **338**, 1587–1593 (2012).
184. Trapnell, C., Pachter, L. & Salzberg, S. L. TopHat: discovering splice junctions with RNA-seq. *Bioinformatics* **25**, 1105–1111 (2009).
185. Steijger, T. *et al.* Assessment of transcript reconstruction methods for RNA-seq. *Nat. Methods* **10**, 1177–1184 (2013).
186. Trapnell, C. *et al.* Transcript assembly and quantification by RNA-seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.* **28**, 511–515 (2010).
187. Djebali, S. *et al.* Landscape of transcription in human cells. *Nature* **489**, 101–108 (2012).
188. Dehm, S. M. & Tindall, D. J. Alternatively spliced androgen receptor variants. *Endocr. Relat. Cancer* **18**, 96 (2011).
189. Antonarakis, E. S. *et al.* AR-V7 and resistance to enzalutamide and abiraterone in prostate cancer. *N. Engl. J. Med.* **371**, 1028–1038 (2014).
190. Xing, Y. & Lee, C. J. Protein modularity of alternatively spliced exons is associated with tissue-specific regulation of alternative splicing. *PLoS Genet.* **1**, e34 (2005).
191. Yeo, G. W., Van Nostrand, E., Holste, D., Poggio, T. & Burge, C. B. Identification and analysis of alternative splicing events conserved in human and mouse. *Proc. Natl Acad. Sci. USA* **102**, 2850–2855 (2005).
192. Merkin, J., Russell, C., Chen, P. & Burge, C. B. Evolutionary dynamics of gene and isoform regulation in mammalian tissues. *Science* **338**, 1593–1599 (2012).
193. Odom, D. T. *et al.* Tissue-specific transcriptional regulation has diverged significantly between human and mouse. *Nat. Genet.* **39**, 730–732 (2007).
194. Schmidt, D. *et al.* Five-vertebrate ChIP-seq reveals the evolutionary dynamics of transcription factor binding. *Science* **328**, 1036–1040 (2010).
195. Moran-Jones, K., Grindlay, J., Jones, M., Smith, R. & Norman, J. C. hnRNP A2 regulates alternative mRNA splicing of *TP53/INP2* to control invasive cell migration. *Cancer Res.* **69**, 9219–9227 (2009).
196. LeFave, C. V., Squatrito, M. & Vorlova, S. Splicing factor hnRNPH drives an oncogenic splicing switch in gliomas. *EMBO J.* **30**, 4084–4097 (2011).
197. Xu, Y. *et al.* Cell type-restricted activity of hnRNPM promotes breast cancer metastasis via regulating alternative splicing. *Genes Dev.* **28**, 1191–1203 (2014).
198. Adler, A. S. *et al.* An integrative analysis of colon cancer identifies an essential function for PRPF6 in tumor growth. *Genes Dev.* **28**, 1068–1084 (2014).
199. Izaguirre, D. I. *et al.* PTBP1-dependent regulation of USP5 alternative RNA splicing plays a role in glioblastoma tumorigenesis. *Mol. Carcinog.* **51**, 895–906 (2012).
200. Fushimi, K. *et al.* Up-regulation of the proapoptotic caspase 2 splicing isoform by a candidate tumor suppressor, RBM5. *Proc. Natl Acad. Sci. USA* **105**, 15708–15713 (2008).
201. Bonnal, S. *et al.* RBM5/Luca-15/H37 regulates Fas alternative splice site pairing after exon definition. *Mol. Cell* **32**, 81–95 (2008).
202. Zhou, X. *et al.* BCLAF1 and its splicing regulator SRSF10 regulate the tumorigenic potential of colon cancer cells. *Nat. Commun.* **5**, 4581 (2014).
203. Wu, S.-J. J. *et al.* The clinical implication of SRSF2 mutation in patients with myelodysplastic syndrome and its stability during disease evolution. *Blood* **120**, 3106–3111 (2012).
204. Damm, F. *et al.* Mutations affecting mRNA splicing define distinct clinical phenotypes and correlate with patient outcome in myelodysplastic syndromes. *Blood* **119**, 3211–3218 (2012).

Acknowledgements

H.D. is supported by a grant from the US Department of Defense Breast Cancer Research Program (W81XWH-14-1-0044). E.K. is supported by the Worldwide Cancer Research Fund. R.K.B. and O.A.-W. are supported by grants from the Edward P. Evans Foundation, the Department of Defense Bone Marrow Failure Research Program (BM150092) and National Institutes of Health/National Heart, Lung and Blood Institute (NIH/NHLBI) (R01 HL128239). O.A.-W. is supported by an NIH K08 Clinical Investigator Award (1K08CA160647-01), a US Department of Defense Postdoctoral Fellow Award in Bone Marrow Failure Research (W81XWH-12-1-0041), the Starr Cancer Consortium (18-A8-075), the Josie Robertson Investigator Program, a Damon Runyon Clinical Investigator Award with support from the Evans Foundation, the Mr William H. Goodwin and Mrs Alice Goodwin Commonwealth Foundation for Cancer Research, and the Experimental Therapeutics Center of Memorial Sloan Kettering Cancer Center. R.K.B. is supported by the Ellison Medical Foundation (AG-NS-1030-13) and NIH/National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) (R01 DK103854).

Competing interests statement

The authors declare no competing interests.